

Regular Kac-Moody superalgebras and integrable highest weight modules

Crystal Hoyt^{*†}

October 15, 2008

Abstract

We define regular Kac-Moody superalgebras and classify them using integrable modules. We give conditions for irreducible highest weight modules of regular Kac-Moody superalgebras to be integrable. This paper is a major part of the proof for the classification of finite-growth contragredient Lie superalgebras.

Keywords: Lie superalgebra; integrable modules; odd reflections.

0 Introduction

The results of this paper are a crucial part of the proof for the classification of contragredient Lie superalgebras with finite growth, and in particular, for the classification of finite-growth Kac-Moody superalgebras [3, 4]. Previously, such a classification was known only for contragredient Lie superalgebras with either symmetrizable Cartan matrices [12, 13], or Cartan matrices with no zeros on the main diagonal, i.e. contragredient Lie superalgebras without simple isotropic roots [6]. Several of the results of this paper are surveyed in [11].

A contragredient Lie superalgebra $\mathfrak{g}(A)$ is a Lie superalgebra defined by a Cartan matrix A [5, 7]. A Lie superalgebra usually has more than one Cartan matrix. However, an odd reflection at a regular simple isotropic root allows one to move from one base to another [1, 10]. An odd reflection yields a new Cartan matrix A' such that $\mathfrak{g}(A')$ and $\mathfrak{g}(A)$ are isomorphic as Lie superalgebras.

A matrix which satisfies certain numerical conditions is called a generalized Cartan matrix (see Definition 1.7). If A is a generalized Cartan matrix then all simple isotropic roots are regular. A contragredient Lie superalgebra $\mathfrak{g}(A)$ is said to be regular Kac-Moody if A and any matrix A' , obtained by a sequence of odd reflections of A , are generalized Cartan matrices. If A is a generalized Cartan matrix and $\mathfrak{g}(A)$ has no simple isotropic roots, then $\mathfrak{g}(A)$ is regular Kac-Moody by definition. Hence, we restrict our attention to regular Kac-Moody

^{*}Address: Crystal Hoyt, Department of Mathematics, Weizmann Institute of Science, Rehovot 76100 Israel; Fax: 972-8-934-6023; Email: crystal.hoyt@weizmann.ac.il

[†]Supported in part by NSF grant DMS-0354321

superalgebras which have a simple isotropic root. Remarkably, there are only a finite number of such families.

It is shown in [4] that if $\mathfrak{g}(A)$ is a finite-growth contragredient Lie superalgebra and the defining matrix A has no zero rows, then simple root vectors of $\mathfrak{g}(A)$ act locally nilpotently on the adjoint module. This implies certain conditions on A which are only slightly weaker than the conditions for the matrix to be a generalized Cartan matrix. For a finite-growth Lie superalgebra, these matrix conditions should still hold after odd reflections, which leads to the definition of a regular Kac-Moody superalgebra. Remarkably, these superalgebras almost always have finite growth. The exception is the family: $Q^\pm(m, n, t)$ with $m, n, t \in \mathbb{Z}_{\geq 1}$ and not all equal to -1 .

By comparing the classification of regular Kac-Moody superalgebras given in this paper to the classification of symmetrizable finite-growth contragredient Lie superalgebras in [12, 13] we obtain the following formulation of our classification theorem.

Theorem 2.26. *If A is a symmetrizable indecomposable matrix and the contragredient Lie superalgebra $\mathfrak{g}(A)$ has a simple isotropic root, then $\mathfrak{g}(A)$ is a regular Kac-Moody superalgebra if and only if it has finite growth.*

If A is a non-symmetrizable indecomposable matrix and the contragredient Lie superalgebra $\mathfrak{g}(A)$ has a simple isotropic root, then $\mathfrak{g}(A)$ is a regular Kac-Moody superalgebra if and only if it belongs to one of the following three families: $q(n)^{(2)}$, $S(1, 2, \alpha)$ with $\alpha \in \mathbb{C} \setminus \mathbb{Z}$, $Q^\pm(m, n, t)$ with $m, n, t \in \mathbb{Z}_{\leq -1}$.

The non-symmetrizable contragredient Lie superalgebra $S(1, 2, \alpha)$ appears in the list of conformal superalgebras [8]. It has finite growth, but is not regular Kac-Moody when $\alpha \in \mathbb{Z}$. The non-symmetrizable regular Kac-Moody superalgebra $Q^\pm(m, n, t)$ was discovered during this classification (see Section 3). If $m, n, t = -1$ then $Q^\pm(m, n, t)$ is just $q(3)^2$, which has finite growth. Otherwise, $Q^\pm(m, n, t)$ has infinite growth, but is hyperbolic for small (absolute) values of m, n , and t . An explicit realization of $Q^\pm(m, n, t)$ is still unknown and would be interesting.

For the proof of Theorem 2.26, we classify the corresponding connected regular Kac-Moody diagrams (see Section 1.3) by using induction on the number of vertices (i.e. simple roots). A subdiagram of a regular Kac-Moody diagram is regular Kac-Moody, however if the subdiagram does not have an isotropic vertex then it is not part of the classification. We work around this difficulty by using odd reflections.

We say that a regular Kac-Moody diagram is *subfinite* if it is connected, has an isotropic vertex, and satisfies the condition that all connected proper subdiagrams which have an isotropic vertex are of finite type. In Section 2, we find all subfinite regular Kac-Moody diagrams by extending connected finite type diagrams which have an isotropic vertex. A diagram of a finite dimensional or affine Kac-Moody superalgebra is subfinite regular Kac-Moody. In Section 4, we prove that every connected regular Kac-Moody diagram with an isotropic

vertex is subfinite by using integrable modules and some explicit computations.

A weight module V of a regular Kac-Moody superalgebra $\mathfrak{g}(A)$ is called integrable if for each real root α the element $X_\alpha \in \mathfrak{g}(A)_\alpha$ is locally nilpotent on V . This is consistent with the original definition when $\mathfrak{g}(A)$ is an affine Lie superalgebra. It is shown in [9] that most non-twisted affine Lie superalgebras do not have non-trivial irreducible integrable highest weight modules. The only exceptions are $osp(1, 2n)^{(1)}$, $A(0, m)^{(1)}$ and $C(n)^{(1)}$. By the same method, one can show that the twisted affine Lie superalgebras, including $q(n)^{(2)}$, but excluding $sl(1, 2n)^{(2)}$, $sl(1, 2n - 1)^{(4)}$ and $osp(2, 2n)^{(2)}$, have only trivial irreducible integrable highest weight modules.

A regular Kac-Moody superalgebra is integrable under the adjoint action, hence, so are its submodules. A non-trivial extension of a regular Kac-Moody diagram Γ yields a non-trivial integrable highest weight module of $\mathfrak{g}(A_\Gamma)$. Thus, if the Kac-Moody superalgebra corresponding to Γ does not have non-trivial irreducible integrable highest weight modules, then Γ is not extendable. Thus, it remains to show that the diagrams for $A(0, m)^{(1)}$, $C(n)^{(1)}$, $S(1, 2, \alpha)$, and $Q^\pm(m, n, t)$ are not extendable, which we do by direct computation.

Integrable irreducible highest weight modules for affine Lie superalgebras were described in [9]. If $L(\lambda)$ is an irreducible highest weight module a regular Kac-Moody superalgebra which does not have an isotropic root, then $L(\lambda)$ is integrable if and only if $\lambda_i \in 2^{p(i)}\mathbb{Z}_{\geq 0}$. We describe the integrable irreducible highest weight modules for the remaining regular Kac-Moody superalgebras which have an isotropic root: $S(1, 2, \alpha)$ with $\alpha \in \mathbb{C} \setminus \mathbb{Z}$, and $Q^\pm(m, n, t)$ with $m, n, t \in \mathbb{Z}_{\geq 1}$ and not all equal to -1 . We show that these superalgebras have non-trivial irreducible integrable highest weight modules, and give explicit conditions on the weights for an irreducible highest weight module to be integrable.

Acknowledgement. The author would like to express her sincere gratitude to Professor Vera Serganova for helpful discussions and suggestions while supervising this dissertation work.

1 Preliminaries

1.1 Contragredient Lie superalgebras

Let A be a $n \times n$ matrix over $\mathbb{C}$, $I = \{1, \dots, n\}$ and $p : I \rightarrow \mathbb{Z}_2$ be a parity function. Fix a vector space $\mathfrak{h}$ over $\mathbb{C}$ of dimension $2n - \text{rk}(A)$. Let $\alpha_1, \dots, \alpha_n \in \mathfrak{h}^*$ and $h_1, \dots, h_n \in \mathfrak{h}$ be linearly independent elements satisfying $\alpha_j(h_i) = a_{ij}$, where a_{ij} is the ij -th entry of A . Define a Lie superalgebra $\bar{\mathfrak{g}}(A)$ by generators $X_1, \dots, X_n, Y_1, \dots, Y_n$ and $\mathfrak{h}$, and by relations

$$[h, X_i] = \alpha_i(h) X_i, \quad [h, Y_i] = -\alpha_i(h) Y_i, \quad [X_i, Y_j] = \delta_{ij} h_i, \quad (1)$$

where the parity of X_i and Y_i is $p(i)$, and the elements of $\mathfrak{h}$ are even.

The *contragredient Lie superalgebra* given by the matrix A is defined to be the quotient of $\bar{\mathfrak{g}}(A)$ by the unique maximal submodule that intersects $\mathfrak{h}$ trivially, and it is denoted by $\mathfrak{g}(A)$. We call A the *Cartan matrix* of $\mathfrak{g}(A)$. If $B = DA$ for some invertible diagonal matrix D , then $\mathfrak{g}(B) \cong \mathfrak{g}(A)$. Hence, we may assume without loss of generality that $a_{ii} \in \{0, 2\}$ for $i \in I$.

The matrix A is said to be *symmetrizable* if there exists an invertible diagonal matrix D such that $B = DA$ is a symmetric matrix, i.e. $b_{ij} = b_{ji}$ for all $i, j \in I$. In this case, we also say that $\mathfrak{g}(A)$ is symmetrizable. The matrix A is *indecomposable* if the set I can not be decomposed into the disjoint union of non-empty subsets J, K such that $a_{j,k} = a_{k,j} = 0$ whenever $j \in J$ and $k \in K$. The proofs for the following lemmas can be found in [4].

Lemma 1.1. *For any subset $J \subset I$ the subalgebra $\mathfrak{a}_J$ in $\mathfrak{g}(A)$ generated by $\mathfrak{h}$, X_i and Y_i , with $i \in J$, is isomorphic to $\mathfrak{h}' \oplus \mathfrak{g}(A_J)$, where A_J is the submatrix of A with coefficients $(a_{ij})_{i,j \in J}$ and $\mathfrak{h}'$ is a subspace of $\mathfrak{h}$. More precisely, $\mathfrak{h}'$ is a maximal subspace in $\cap_{i \in J} \text{Ker } \alpha_i$ which trivially intersects the span of h_i , $i \in J$.*

A superalgebra $\mathfrak{g} = \mathfrak{g}(A)$ has a natural $\mathbb{Z}$ -grading $\mathfrak{g} = \oplus \mathfrak{g}_m$, called the *principal grading*, which is defined by $\mathfrak{g}_0 := \mathfrak{h}$ and $\mathfrak{g}_1 := \mathfrak{g}_{\alpha_1} \oplus \cdots \oplus \mathfrak{g}_{\alpha_n}$. We say that $\mathfrak{g}$ is of *finite growth* if $\dim \mathfrak{g}_n$ grows polynomially depending on n . This means that the Gelfand-Kirillov dimension of $\mathfrak{g}$ is finite. We recall some results from [4].

Theorem 1.2 (Hoyt, Serganova). *Suppose A is a matrix with no zero rows. If $\mathfrak{g}(A)$ has finite growth, then ad_{X_i} are locally nilpotent for all $i \in I$.*

Lemma 1.3. *Let $\mathfrak{g}(A)$ be a contragredient Lie superalgebra. Then ad_{X_i} are locally nilpotent for all $i \in I$ if and only if A satisfies the following conditions, (after rescaling the rows of A such that $a_{ii} \in \{0, 2\}$ for all $i \in I$):*

1. *if $a_{ii} = 0$ and $p(i) = 0$, then $a_{ij} = 0$ for all $j \in I$;*
2. *if $a_{ii} = 2$, then $a_{ij} \in 2^{p(i)}\mathbb{Z}_{\leq 0}$ for all $j \in I$;*
3. *if $a_{ij} = 0$ and $a_{ji} \neq 0$, then $a_{ii} = 0$.*

1.2 Roots and Reflections

The Lie superalgebra $\mathfrak{g} = \mathfrak{g}(A)$ has a *root space decomposition*

$$\mathfrak{g} = \mathfrak{h} \oplus \bigoplus_{\alpha \in \Delta} \mathfrak{g}_{\alpha}.$$

Every root is either a positive or a negative linear combination of the simple roots, $\alpha_1, \dots, \alpha_n$. Accordingly, we have the decomposition $\Delta = \Delta^+ \cup \Delta^-$, and we call $\alpha \in \Delta^+$ positive and $\alpha \in \Delta^-$ negative. One can define $p : \Delta \rightarrow \mathbb{Z}_2$ by letting $p(\alpha) = 0$ or 1 whenever α is even or odd, respectively. By Δ_0 (resp. Δ_1) we denote the set of even (resp. odd) roots.

Let $\Pi := \{\alpha_1, \dots, \alpha_n\}$ be the set of simple roots of $\mathfrak{g}(A)$. There are four possibilities for each simple root α_i :

1. if $a_{ii} = 2$ and $p(\alpha_i) = 0$, then X_i, Y_i and h_i generate a subalgebra isomorphic to $sl(2)$;
2. if $a_{ii} = 0$ and $p(\alpha_i) = 0$, then X_i, Y_i and h_i generate a subalgebra isomorphic to the Heisenberg algebra;
3. if $a_{ii} = 2$ and $p(\alpha_i) = 1$, then X_i, Y_i and h_i generate a subalgebra isomorphic to $osp(1|2)$, and in this case $2\alpha_i \in \Delta$;
4. if $a_{ii} = 0$ and $p(\alpha_i) = 1$, then X_i, Y_i and h_i generate a subalgebra isomorphic to $sl(1|1)$.

In the last case we say that α_i is *isotropic*, and otherwise α_i is *non-isotropic*. A simple root α_i is *regular* if for any other simple root α_j , $a_{ij} = 0$ implies $a_{ji} = 0$.

If α_k is a simple root with $a_{kk} \neq 0$, we define the (even) reflection r_k at α_k by

$$r_k(\alpha_i) = \alpha_i - \alpha_i(h_k)\alpha_k, \quad \alpha_i \in \Pi.$$

If α_k is a regular isotropic root, we define the *odd reflection* r_k at α_k as follows:

$$r_k(\alpha_i) = \begin{cases} -\alpha_k, & \text{if } i = k; \\ \alpha_i, & \text{if } a_{ik} = a_{ki} = 0, i \neq k; \\ \alpha_i + \alpha_k, & \text{if } a_{ik} \neq 0 \text{ or } a_{ki} \neq 0, i \neq k; \end{cases}$$

$$X'_i := \begin{cases} Y_i, & \text{if } i = k; \\ X_i, & \text{if } i \neq k, \text{ and } a_{ik} = a_{ki} = 0; \\ [X_i, X_k], & \text{if } i \neq k, \text{ and } a_{ik} \neq 0 \text{ or } a_{ki} \neq 0; \end{cases}$$

$$Y'_i := \begin{cases} X_i, & \text{if } i = k; \\ Y_i, & \text{if } i \neq k, \text{ and } a_{ik} = a_{ki} = 0; \\ [Y_i, Y_k], & \text{if } i \neq k, \text{ and } a_{ik} \neq 0 \text{ or } a_{ki} \neq 0; \end{cases}$$

and

$$h'_i := [X'_i, Y'_i].$$

Then

$$h'_i = \begin{cases} (-1)^{p(\alpha_i)}(a_{ik}h_k + a_{ki}h_i) & \text{if } i \neq k, \text{ and } a_{ik} \text{ or } a_{ki} \neq 0, \\ h_i & \text{if } i \neq k, \text{ and } a_{ik} = a_{ki} = 0, \\ h_k & \text{if } i = k. \end{cases}$$

Set $\alpha'_i := r_k(\alpha_i)$ for $i \in I$.

Lemma 1.4. *The roots $\alpha'_1, \dots, \alpha'_n$ are linearly independent. The elements $X'_1, \dots, X'_n, Y'_1, \dots, Y'_n$ together with $h'_1, \dots, h'_n$ satisfy (1). Moreover, $\mathfrak{h}$ and $X'_1, \dots, X'_n, Y'_1, \dots, Y'_n$ generate $\mathfrak{g}$.*

It follows that given a matrix A and a regular isotropic root α_k , one can construct a new matrix A' such that $\mathfrak{g}(A') \cong \mathfrak{g}(A)$ as superalgebras. Explicitly, the entries of A' are defined by the following (where we assume $i \neq k$ and $j \neq k$):

$$a'_{kk} := a_{kk}; \quad a'_{kj} := a_{kj}; \quad a'_{ik} := -a_{ki}a_{ik};$$

$$a'_{ij} := \begin{cases} a_{ij}, & \text{if } a_{ik} = a_{ki} = 0; \\ a_{ki}a_{ij}, & \text{if } a_{ik} \text{ or } a_{ki} \neq 0, \text{ and } a_{kj} = a_{jk} = 0; \\ a_{ki}a_{ij} + a_{ik}a_{kj} + a_{ki}a_{ik}, & \text{if } a_{ik} \text{ or } a_{ki} \neq 0, \text{ and } a_{jk} \text{ or } a_{kj} \neq 0. \end{cases}$$

We say that A' is obtained from A (and $\Pi' := \{\alpha'_1, \dots, \alpha'_n\}$ is obtained from Π) by an odd reflection with respect to α_k . If Δ'^+ is the set of positive roots with respect to Π' , then

$$\Delta'^+ = (\Delta^+ \setminus \{\alpha_k\}) \cup \{-\alpha_k\}.$$

Remark 1.5. A reflection with respect to a regular isotropic simple root α_k is indeed a reflection. If A'' is obtained from A by successively applying the reflection at α_k twice, then there is an invertible diagonal matrix D such that $A'' = DA$ and scalars b_i, c_i such that $X''_i = b_i X_i$ and $Y''_i = c_i Y_i$. It is possible to define an “odd reflection” at a simple isotropic root which is not regular, but in this case the subalgebra generated by $X'_1, \dots, X'_n, Y'_1, \dots, Y'_n$ and $\mathfrak{h}$ is necessarily a proper subalgebra of $\mathfrak{g}(A)$.

A root $\alpha \in \Delta$ is called *real* if α or $\frac{1}{2}\alpha$ is simple in some base Π' obtained from Π by a sequence of even and/or odd reflections, and it is called *imaginary* otherwise.

Lemma 1.6. *Let Π' be obtained from Π by an odd reflection. Then $\dim \mathfrak{g}'_n \leq \dim \mathfrak{g}_{-2n} \oplus \dots \oplus \mathfrak{g}_{2n}$. In particular, if $\mathfrak{g}(A)$ is of finite growth, then $\mathfrak{g}(A')$ is of finite growth.*

Hence, the notion of finite growth does not depend on the choice of the base for $\mathfrak{g}(A)$. It thus follows from Theorem 1.2 that if $\mathfrak{g}(A)$ has finite growth and A has no zero rows, then A and any matrix A' obtained from A by a sequence of odd reflections satisfies the matrix conditions of Lemma 1.3.

Definition 1.7. A matrix A is called a *generalized Cartan matrix* if it satisfies the matrix conditions of Lemma 1.3, where the third condition is strengthened to the following:

$$3'. \text{ if } a_{ij} = 0, \text{ then } a_{ji} = 0.$$

If a matrix satisfies the conditions of Lemma 1.3, then condition (3') is equivalent to the condition that all simple isotropic roots are regular. We call $\mathfrak{g}(A)$ a *regular Kac-Moody superalgebra* if A and any matrix obtained from A by a sequence of odd reflections is a generalized Cartan matrix.

We call a root $\alpha \in \Delta$ *principal* if either α is even and belongs to some base Π' obtained from Π by a sequence of odd reflections, or if $\alpha = 2\beta$ where β is odd and belongs to some base Π' obtained from Π by a sequence of odd reflections. For a principal root, the subalgebra generated by X_α, Y_α and $h_\alpha := [X_\alpha, Y_\alpha]$ is isomorphic to $\mathfrak{sl}_2$, and we may choose X_α, Y_α such that $\alpha(h_\alpha) = 2$. Note that if $\alpha = 2\beta$, then $X_\alpha = [X_\beta, X_\beta]$.

Let $\Pi_0 \subset \Delta$ denote the set of principal roots. It is clear that $\Pi_0 \subset \Delta_0^+$. In general this set can be infinite, but this is not the case whenever $\mathfrak{g}(A)$ has finite growth [4]. For any finite subset $S \subset \Pi_0$, we can define a matrix B by setting $b_{ij} = \alpha_j(h_i)$.

Lemma 1.8. *If a Lie superalgebra $\mathfrak{g}(A)$ has finite growth, then for any finite subset S of Π_0 the Lie algebra $\mathfrak{g}(B)$ also has finite growth. In particular, B is a generalized Cartan matrix for a finite growth Kac-Moody algebra.*

1.3 Matrix diagrams

Given a Cartan matrix A , we can associate a *matrix diagram*, denoted Γ_A (or simply Γ when A is fixed), as follows. After possibly rescaling A , we may assume that $a_{ii} = 0$ or 2 . The vertices of Γ_A correspond to the simple roots of $\mathfrak{g}(A)$ and are given by the following table.

$\mathfrak{g}(A)$	A	$p(1)$	Diagram
A_1	(2)	0	$\circ$
$B(0, 1)$	(2)	1	$\bullet$
$A(0, 0)$	(0)	1	$\otimes$
Heisenberg	(0)	0	$\square$

We join vertex v_i to vertex v_j by an arrow if $a_{ij} \neq 0$, and we label this arrow with the number a_{ij} . This correspondence between Cartan matrices and matrix diagrams is a bijection. The condition that the matrix A is indecomposable corresponds to the requirement that the diagram Γ_A is connected. We say that a vertex is isotropic if it is of the type $\otimes$. We use the notation of the following table to denote the possible vertex types.

Notation	Vertex Types
$\bullet$	$\circ$ or $\otimes$
$\odot$	$\circ$ or $\bullet$
$\textcircled{v}$	$\circ$ or $\otimes$ or $\bullet$

Let A be a Cartan matrix with $a_{ii} = 0$. Then rescaling the i^{th} row of the matrix A (i.e. multiplying by a non-zero constant) corresponds to rescaling all labels of arrows exiting the vertex v_i of the diagram Γ_A . This defines an equivalence relation on matrix diagrams, where $\Gamma_{A'} \sim \Gamma_A$ if $A' = DA$ for some invertible diagonal matrix D , and in this case $\mathfrak{g}(A') \cong \mathfrak{g}(A)$. We consider matrix diagrams modulo this equivalence.

A diagram Γ_A is called a *regular Kac-Moody* (or regular Kac-Moody diagram) if the corresponding contragredient Lie superalgebra $\mathfrak{g}(A)$ is regular Kac-Moody. An odd reflection of a matrix diagram Γ_A at an isotropic vertex v_i is defined to be the diagram $\Gamma_{A'}$, where A' is obtained from A by an odd reflection at the corresponding isotropic root α_i of $\mathfrak{g}(A)$. Note that the set of all regular Kac-Moody diagrams is closed under odd reflections. Denote by $C(\Gamma)$ the collection of all diagrams obtained from sequences of odd reflections of a regular Kac-Moody diagram Γ . When α_i is a simple odd isotropic root, we denote by r_i the odd reflection with respect to α_i .

By a *subdiagram* Γ' of the diagram Γ , we mean a full subdiagram, i.e. if the vertices v_i and v_j are in Γ' then the arrows with labels a_{ij} and a_{ji} also belong to

Γ' . We say that a connected regular Kac-Moody diagram is *extendable* if it is a proper subdiagram of a connected regular Kac-Moody diagram. For a 3-vertex subdiagram $\Gamma' = \{v_i, v_j, v_k\}$, with v_j isotropic, we refer to the fractions $\frac{a_{ji}}{a_{jk}}$ and $\frac{a_{jk}}{a_{ji}}$ as the *ratios* of the isotropic vertex v_j in Γ' .

2 Classification of connected subfinite regular Kac-Moody diagrams

In order to classify regular Kac-Moody superalgebras, we classify the corresponding regular Kac-Moody diagrams. A diagram for a finite dimensional or affine Kac-Moody superalgebra is regular Kac-Moody. We call a regular Kac-Moody diagram *subfinite* if it is connected, it contains an isotropic vertex, and it satisfies the following condition for all reflected diagrams: all connected proper subdiagrams which contain an isotropic vertex are of finite type. Later in Section 5, we will show that all regular Kac-Moody diagrams are subfinite.

In this section, we classify subfinite regular Kac-Moody diagrams, by using induction on the number of vertices. A subdiagram of a subfinite regular Kac-Moody diagram is regular Kac-Moody, however if it does not contain an isotropic vertex then it is not part of our classification. We will work around this difficulty by using odd reflections.

Note that we only need to find the extensions of one diagram Γ belonging to a collection of reflected diagrams $\mathcal{C}(\Gamma)$, and then include all reflections of each extended diagram in our classification.

2.1 Regular Kac-Moody diagrams: 2 or 3 vertices

In this section, we find all connected diagrams with two or three vertices which are regular Kac-Moody and contain an isotropic vertex.

Lemma 2.1. *The connected regular Kac-Moody 2-vertex diagrams which contain an isotropic vertex are $A(1,0)$ and $B(1,1)$.*

Proof.

◊ If $a = -1$, then by reflecting at v_1 we have $A(1,0)$.

$$\otimes \xrightleftharpoons[-1]{1} \circ \quad \xrightarrow{r_1} \quad \otimes \xrightleftharpoons[1]{1} \otimes$$

◊ If $a \neq -1$, then reflecting at v_1 , we have $a, \frac{-a}{a+1} \in \mathbb{Z}_{<0}$ which implies $a = -2$. This is $B(1,1)$.

$$\otimes \xrightleftharpoons[a]{1} \odot \quad \xrightarrow{r_1} \quad \otimes \xrightleftharpoons[\frac{-a}{1+a}]{1} \odot$$

□

We note that all 2-vertex diagrams of regular Kac-Moody superalgebras are of finite type.


Lemma 2.2. *The regular Kac-Moody extensions of $A(1,0)$ to three vertices are the following: $A(0,2)$, $A(1,1)$, $B(1,2)$, $B(2,1)$, $C(3)$, $G(3)$, $D(2,1,\alpha)$, $A(1,2)^{(2)}$, $A(0,1)^{(1)}$, $B(1,1)^{(1)}$, $S(1,2,\alpha)$, $q(3)^{(2)}$, $Q^\pm(m,n,t)$.*

Proof. We consider each case for attaching a vertex to an $A(1,0)$ diagram.


Case 1:

$$\otimes_1 \xrightleftharpoons[1]{1} \otimes_2 \xrightleftharpoons[b]{a} \odot_3 \quad a, b \neq 0$$

◊If v_3 is $\otimes$, then by reflecting at v_2 we have $1+a, 1+\frac{1}{a} \in \mathbb{Z}_{\leq 0}$, which implies $a = -1$. This is $A(1,1)$.




◊If v_3 is $\odot$ and $b = -2$, then by reflecting at v_2 we have $1+a, 2+\frac{2}{a} \in \mathbb{Z}_{\leq 0}$, which implies $a = -1$. This is $B(1,2)$.




◊If v_3 is $\bigcirc$ and $a, b = -1$, then this is $A(0,2)$.


◊If v_3 is $\bigcirc$, $b = -1$ and $a \neq -1$, then by reflecting at v_2 we have $1+a \in \{-1, -2\}$, which implies $a \in \{-2, -3\}$. If $a = -2$ then this is $C(3)$, and if $a = -3$ then this is $G(3)$.



Case 2:



◊ If v_2 is $\otimes$, then after rescaling we have




Then $1 + a + \frac{1}{b}, 1 + b + \frac{1}{c} \in \mathbb{Z}_{<0}$ or both zero. By symmetry it follows that

$$\begin{aligned} 1 + a + \frac{1}{b} &= m \\ 1 + b + \frac{1}{c} &= n \\ 1 + c + \frac{1}{a} &= t \end{aligned} \in \mathbb{Z}_{<0}, \text{ or all equal to zero.} \quad (2)$$

If they all equal zero, then this is $D(2, 1, \alpha)$. If they all equal -1 , then $a, b, c = -1$ and this is $q(3)^{(2)}$. If they are in $\mathbb{Z}_{<0}$ and not all equal to -1 , then this is $Q^\pm(m, n, t)$.

◊ If v_2 is $\odot$ and $b, d = -2$, then by reflecting at v_1 we have $4 + \frac{2}{a}, 4 + \frac{2}{c}, 1 + a + c \in \mathbb{Z}_{<0}$ or all zero, which implies $a, c = -\frac{1}{2}$. If v_2 is $\circ$ then this is $A(1, 2)^{(2)}$, and if v_2 is $\bullet$ then this is $B(1, 1)^{(1)}$.



◊If v_2 is $\bigcirc$, $b = -2$ and $d = -1$, then by reflecting at v_1 we have

$$\begin{array}{ccc} \begin{array}{c} \bigcirc \\ \swarrow a \quad \searrow -1 \\ \bigotimes \quad \bigotimes \\ \xleftarrow{1} \quad \xrightarrow{1} \end{array} & \xrightarrow{\vec{r}_1} & \begin{array}{c} \bullet \\ \swarrow a \quad \searrow 3+\frac{2}{a} \\ \bigotimes \quad \bigcirc \\ \xleftarrow{1} \quad \xrightarrow{-1} \end{array} \end{array} \quad Q = 1 + a + c. \quad (3)$$

Then either $1 + a + c \in \mathbb{Z}_{<0}$ and $3 + \frac{2}{a} \in 2\mathbb{Z}_{<0}$, or both equal zero. If they are both zero, then $a = -\frac{2}{3}, c = -\frac{1}{3}$ and this is $G(3)$.

So now assume that $1 + a + c \neq 0$. Then by reflecting at v_3 we have

$$\begin{array}{ccc} \begin{array}{c} \bigcirc \\ \swarrow a \quad \searrow -1 \\ \bigotimes \quad \bigotimes \\ \xleftarrow{1} \quad \xrightarrow{1} \end{array} & \xrightarrow{\vec{r}_3} & \begin{array}{c} \bigotimes \\ \swarrow 1+a+c \quad \searrow c \\ \bigcirc \quad \bigotimes \\ \xleftarrow{1} \quad \xrightarrow{-1} \end{array} \end{array} \quad (4)$$

This implies $1 + a + c \in \{-1, -2\}$ and $c \neq -\frac{1}{3}$.

Suppose $1 + a + c = -1$. Then $a = -2 - c$ and $c \neq -2$. By reflecting the last diagram of (4) at v_2 we have:

$$\begin{array}{ccc} \begin{array}{c} \bigotimes \\ \swarrow -1-3c \quad \searrow c \\ \bigotimes \quad \bigcirc \\ \xleftarrow{-2} \quad \xrightarrow{2+5c} \end{array} & \xrightarrow{\vec{r}_1} & \begin{array}{c} \bigcirc \\ \swarrow -1-3c \quad \searrow -1 \\ \bigotimes \quad \bullet \\ \xleftarrow{-2} \quad \xrightarrow{2+5c} \end{array} \end{array} \quad \begin{array}{l} R = \frac{4+9c}{2+5c} \\ P = -1 - 3c \end{array}$$

Then, $1 + 3c \neq 0$, $2 + 5c \neq 0$, and $R = \frac{4+9c}{2+5c} \in 2\mathbb{Z}_{<0}$. Hence, c is rational. From (3) we have that $\frac{4+3c}{2+c} = 3 + \frac{2}{a} \in 2\mathbb{Z}_{<0}$. But the system of equations

$$\frac{4+3c}{2+c} \leq -2, \quad \frac{4+9c}{2+5c} \leq -2$$

has no real solution. This is a contradiction.

Suppose $1 + a + c = -2$. Then $a = -3 - c$ and $c \neq -3$. By reflecting the last diagram of (4) at v_2 we have


$$\begin{array}{ccc} \begin{array}{c} \bigotimes \\ \swarrow -2 \quad \searrow c \\ \bigcirc \quad \bigotimes \\ \xleftarrow{1} \quad \xrightarrow{-1} \end{array} & \xrightarrow{\vec{r}_2} & \begin{array}{c} \bigotimes \\ \swarrow -2 \quad \searrow c \\ \bullet \quad \bigcirc \\ \xleftarrow{-2} \quad \xrightarrow{S} \end{array} \end{array} \quad \begin{array}{l} S = \frac{3+7c}{1+3c} \\ P = -1 - 3c \end{array}$$

It follows that $1 + 3c \neq 0$, and $S = \frac{3+7c}{1+3c} \in 2\mathbb{Z}_{<0}$. Hence, c is rational. From (3) we have that $\frac{7+3c}{3+c} = 3 + \frac{2}{a} \in 2\mathbb{Z}_{<0}$. But, the system of equations

$$\frac{7+3c}{3+c} \leq -2, \quad \frac{3+7c}{1+3c} \leq -2$$


has no real solution. This is a contradiction.

◊If v_2 is $\bigcirc$ and $b, d = -1$, then by reflecting at v_1 we have




Then either $1 + a + c = 0$ and $1 + 2a = 0$, or $1 + a + c \in \{-1, -2\}$. But $1 + a + c \neq -2$ by the previous case, since this is not a diagram for $G(3)$. If $1 + a + c = 0$, then this is $C(3)$.


If $1 + a + c = -1$, then $a = -2 - c$ and $c \neq -2$. If $c = -1$, then this is $A(0, 1)^{(1)}$. For $c \neq -1$ the substitution $c = \frac{1-\alpha}{\alpha}$ is reversible. After substituting and rescaling, we have



This diagram is $S(1, 2, \alpha)$, and is regular Kac-Moody precisely when α is not an integer. Indeed, by reflecting the following diagram at v_1 we have:



Reflecting at v_3 gives us



By induction, two diagrams given by labels α_1 and α_2 are connected by a sequence of odd reflections precisely when $\alpha_1 - \alpha_2 \in \mathbb{Z}$. Hence, $S(1, 2, \alpha)$ is regular Kac-Moody if and only if α is not an integer.

We have found all regular Kac-Moody extensions of $A(1, 0)$ by one vertex. $\square$

Lemma 2.3. *The regular Kac-Moody extensions of $B(1, 1)$ to three vertices that are not extensions of $A(1, 0)$ are the following: $A(2, 2)^{(4)}$, $D(2, 1)^{(2)}$.*

Proof. We consider each case for attaching a vertex to an $B(1, 1)$ diagram.

Case 1:

$$\otimes_1 \xrightleftharpoons[-2]{1} \bullet_2 \xrightleftharpoons[b]{a} \odot_3 \quad a, b \neq 0$$

Reflecting at v_1 we have


$$\otimes \xrightleftharpoons[-2]{1} \bullet \xrightleftharpoons[b]{a} \odot \xrightarrow{r_1} \otimes \xrightleftharpoons[-2]{1} \circ \xrightleftharpoons[b]{-a} \odot .$$

This implies $a, -a \in \mathbb{Z}_{<0}$, which is a contradiction.

Case 2:


$$\odot_1 \xrightleftharpoons[a]{-2} \otimes_2 \xrightleftharpoons[-2]{1} \bullet_3 \quad a \neq 0$$

By reflecting at v_2 we have




which implies $2 + 2a, 2 + \frac{2}{a} \in \mathbb{Z}_{<0}$. The unique solution is $a = -1$. If v_1 is $\odot$ then this is $A(2, 2)^{(4)}$, and if v_1 is $\bullet$ then this is $D(2, 1)^{(2)}$.

Case 3:



Now $a \neq 0$, $b \in \mathbb{Z}_{<0}$, and $c \in 2\mathbb{Z}_{<0}$. By reflecting at v_1 we have



where $P = 2 - b + \frac{2}{a}$, $Q = 2 - c + 2a$, and $P, Q \in \mathbb{Z}_{\leq 0}$. But, the system of equations

$$b \leq -1, \quad c \leq -2, \quad 2 - c + 2a \leq 0, \quad 2 - b + \frac{2}{a} \leq 0$$

has no real solution. This is a contradiction. $\square$

We note that the only regular Kac-Moody diagram with three vertices which is not finite or of finite growth is $Q^\pm(m, n, t)$.

2.2 Preliminaries

We say that an n -vertex diagram is a *cycle* if it is a connected diagram and each vertex is connected to exactly two other vertices. We say that an n -vertex diagram is a *chain* if the vertices can be enumerated by the set $\{1, 2, \dots, n\}$ such that $a_{ij} = 0$ if and only if $j \neq i + 1$ and $i \neq j + 1$. A proper connected subdiagram of a cycle is a chain.

Lemma 2.4. *Suppose that Γ is a subfinite regular Kac-Moody n -vertex chain where the vertex v_n is isotropic and the vertices v_i for $1 < i < n$ are not isotropic. Then there is a sequence of odd reflections of the vertices v_i with $3 \leq i \leq n$ such that in the reflected diagram Γ' the vertex v_2 is isotropic. The vertex v_1 is unchanged and $a'_{12} = a_{12}$.*

Proof. If $n = 2$, then we are done. So suppose the lemma holds for such a chain with $n - 1$ vertices. Let Γ be a chain with n vertices satisfying the hypothesis of the lemma. Now $a_{n,n} = 0$ because v_n is isotropic, and $a_{n-1,n-1} = 2$ because v_{n-1} is not isotropic. Since Γ is a chain, $a_{n-1,n-2} \neq 0$ and $a_{n-2,n} = a_{n,n-2} = 0$. Since Γ is a subfinite regular Kac-Moody diagram, the subdiagram

containing the vertices v_{n-2}, v_{n-1}, v_n is of finite type, which by the three vertex classification implies that $a_{n-1,n} = -1$ and the vertex v_{n-1} is even.

By reflecting at v_n we obtain a diagram Γ' where v'_{n-1} is isotropic. We observe that Γ' is again a chain. Indeed, $a'_{in} = a'_{ni} = 0$ and $a'_{ij} = a_{ij}$ for $i < n-1$, $j < n$, because $a_{in} = a_{ni} = 0$ for $i \neq n-1$. Also, $a'_{n-1,i} = a_{n,n-1}a_{n-1,i} = 0$ for $i < n-2$. We can apply the induction hypothesis to $\Gamma' - \{v_n\}$ which gives us the desired sequence of odd reflections of Γ' and hence of Γ . $\square$

Lemma 2.5. *Let Γ be a subfinite regular Kac-Moody diagram. Suppose Γ' is a connected m -vertex subdiagram of Γ such that Γ' does not contain an isotropic vertex. Then there is a sequence of odd reflections R of Γ such that $R(\Gamma') = \Gamma'$, and there is a connected $(m+1)$ -vertex subfinite regular Kac-Moody subdiagram of $R(\Gamma)$ which contains $\Gamma' = R(\Gamma')$ as a subdiagram and an isotropic vertex.*

Proof. First note that Γ contains an isotropic vertex because it is subfinite. So we may take a minimal subdiagram Γ'' containing Γ' and an isotropic vertex. Then by the minimality of Γ'' the set of vertices in $\Gamma'' - \Gamma'$ form a chain with an isotropic vertex, and only one of these vertices is connected to the subdiagram Γ' . Denote this vertex v_2 and the rest of the vertices in the chain $\Gamma'' - \Gamma'$ by $\{v_3, \dots, v_n\}$ so that v_k is connected to v_{k-1} and v_{k+1} . Then v_n is isotropic by minimality. Choose any vertex in Γ' that is connected to v_2 and denote it v_1 .

Now apply the Lemma 2.4 to the subdiagram given by $\{v_1, v_2, \dots, v_n\}$ to obtain a sequence of odd reflections R such that $R(v_2)$ is isotropic. By minimality of Γ'' each vertex v_k for $k \geq 3$ is not connected to Γ' , so each reflection does not change the subdiagram Γ' , ie. $a'_{ij} = a_{ij}$ if $v_i, v_j \in \Gamma'$. Then the subdiagram $R(\Gamma' \cup \{v_2\})$ of $R(\Gamma)$ is a subfinite regular Kac-Moody $m+1$ -vertex diagram containing the diagram Γ' and an isotropic vertex. $\square$

Lemma 2.6. *If Γ is a subfinite regular Kac-Moody diagram with $n \geq 4$ vertices, then any odd non-isotropic vertex of Γ has degree one.*

Proof. First note that connected finite type 3-vertex regular Kac-Moody diagrams satisfy the condition that odd non-isotropic vertices have degree one. Suppose that the vertex v_2 is odd non-isotropic with degree greater than or equal to two. Let $\Gamma' = \{v_1, v_2, v_3\}$, where $a_{12}, a_{21}, a_{23}, a_{32} \neq 0$. By the 3-vertex classification, Γ' does not contain an isotropic vertex since Γ is subfinite. By Lemma 2.5, we are reduced to the case when Γ has four vertices.

If v_4 is connected to an odd non-isotropic vertex v_i , then v_i has degree two in a 3-vertex subdiagram $\{v_4, v_i, v_j\}$, which contradicts the assumption that Γ is subfinite. Thus, $a_{24} = a_{42} = 0$. We may assume $a_{14}, a_{41} \neq 0$, which implies that v_1 is even. By reflecting at v_4 we obtain a diagram in which $R(v_1)$ is isotropic, $R(v_2) = v_2$ and $a'_{2i} = a_{2i}$ for $i = 1, \dots, 4$. But then the subdiagram $R(\{v_1, v_2, v_3\})$ is not of finite type. This contradicts the assumption that Γ is subfinite. $\square$

Lemma 2.7. *If Γ is a subfinite regular Kac-Moody diagram with $n \geq 4$ vertices, and Γ' is a 3-vertex subdiagram with an isotropic vertex of degree two in Γ'*

such that the ratio of the vertex is not a negative rational number, then Γ' is a $D(2, 1, \alpha)$ diagram.

Proof. Since Γ' is finite type regular Kac-Moody, this follows from the 3-vertex classification. $\square$

Corollary 2.8. *If Γ is a subfinite regular Kac-Moody diagram with $n \geq 4$ vertices and Γ' is a 4-vertex subdiagram containing an isotropic vertex of degree three in Γ' , then Γ' contains a $D(2, 1, \alpha)$ diagram containing this vertex.*

2.3 Subfinite regular Kac-Moody: 4 vertices

We introduce the following notation in order to simplify the presentation. If Γ is a diagram and $\{v_i\}_{i \in I}$ is a subset of the vertices of Γ , then we denote by $\Gamma_{\{i| i \in I\}}$ the subdiagram of Γ obtained by removing the vertices $\{v_i\}_{i \in I}$. For example, $\Gamma_{1,2}$ is the subdiagram of Γ obtained by removing the vertices v_1 and v_2 . Let $\mathcal{F}$ denote the set of all connected finite type regular Kac-Moody diagrams which contain an isotropic vertex.

For each finite type 3-vertex diagram, we will consider each case for attaching an additional vertex to the diagram. Let Γ denote the corresponding extended diagram.

Lemma 2.9. *The subfinite regular Kac-Moody extensions of $D(2, 1, \alpha)$ to four vertices are $D(3, 1)$, $D(2, 2)$, F_4 , $B(2, 1)^{(2)}$, $D(2, 1, \alpha)^{(1)}$, $G(3)^{(1)}$, $A(1, 3)^{(2)}$, and $A(2, 3)^{(2)}$.*

Proof. We consider each case for attaching a vertex to a $D(2, 1, \alpha)$ diagram.

Case 1:

$a, b, c, d, e \neq 0$


$$\begin{aligned} a + \frac{1}{b} &= -1 \\ b + \frac{1}{c} &= -1 \\ c + \frac{1}{a} &= -1 \end{aligned} \quad (5)$$

◊If v_4 is $\otimes$, then $\Gamma_2 \in \mathcal{F}$ implies $d = -1$ and $\Gamma_3 \in \mathcal{F}$ implies $c = -d$. Then by (5) we have $a = \frac{-1}{2}$, $b = -2$, $c = 1$, and $d = -1$. This is $D(2, 2)$.

◊If v_4 is $\odot$ and $e = -2$, then $\Gamma_2 \in \mathcal{F}$ implies $d = -1$ and $\Gamma_3 \in \mathcal{F}$ implies $c = -d$. Then by (5) we have $a = \frac{-1}{2}$, $b = -2$, $c = 1$, and $d = -1$. If v_4 is $\circ$, then this is $B(2, 1)^{(1)}$. If v_4 is $\bullet$, then this is $A(2, 3)^{(2)}$.


◊If v_4 is $\circ$ and $e = -1$, then $\Gamma_2 \in \mathcal{F}$ implies $d \in \{-1, -2, -3\}$ and $\Gamma_3 \in \mathcal{F}$ implies $c \in \{-d, \frac{-d}{2}, \frac{-d}{3}\}$. If $d \in \{-1, -2\}$, then $c \in \{-d, \frac{-d}{2}\}$. This yields three distinct options, and we find that Γ is one of the following: $D(3, 1)$, F_4 ,

$A(1, 3)^{(2)}$. If $d = -3$, then $c \in \{3, \frac{3}{2}, 1\}$ and by reflecting at v_1 we obtain



Then $r_1(\Gamma)_1 \in \mathcal{F}$ implies $\frac{3-c}{2} \leq 0$. Hence, $c = 3$. Then by (5) we have $a = \frac{-1}{4}$, $b = \frac{-4}{3}$. This is $G(3)^{(1)}$.

Case 2:




◊If v_4 is $\otimes$, then $\Gamma_3 \in \mathcal{F}$ implies $d = -c$ and $\Gamma_1 \in \mathcal{F}$ implies $f = -1$. Then by substitution and by (6), we have $d + \frac{f}{a} = -(c + \frac{1}{a}) = 1$. Now Γ_2 must be a $D(2, 1, \alpha)$ diagram, so (2) implies $d + \frac{f}{a} = -1$. This is a contradiction.

◊If v_4 is $\odot$, $e = -1$ and $g = -2$, then $\Gamma_2 \in \mathcal{F}$ implies $a = \frac{3}{2}$, $d = \frac{-1}{2}$. Now $\Gamma_3 \in \mathcal{F}$ implies $\frac{d}{c} < 0$. Hence, $c > 0$. Then $c + \frac{1}{a} > 0$, which contradicts (6).

◊If v_4 is $\odot$ and $e = g = -1$, then $\Gamma_2 \in \mathcal{F}$ implies $d = \frac{f}{a} = \frac{-1}{2}$, $\Gamma_1 \in \mathcal{F}$ implies $f < 0$, and $\Gamma_3 \in \mathcal{F}$ implies $\frac{d}{c} < 0$. Since $d, f < 0$ we have $a, c > 0$. But then $c + \frac{1}{a} > 0$, which contradicts (6).

Case 3:




Suppose v_4 is $\odot$. Then $\frac{d}{b}, \frac{a}{d}, \frac{c}{e}, \frac{e}{b}, \frac{f}{a}, \frac{c}{f} \in \mathbb{Q}_{<0}$ implies $\frac{a+c}{b} > 0$, which contradicts (7). Hence, v_4 is $\otimes$. Thus all subdiagrams are $D(2, 1, \alpha)$ diagrams. So we have $g = d$, $h = e$, and $k = f$. We also have $a + b + c = 0$, $c + d + e = 0$,

$a + e + f = 0$, and $b + d + f = 0$. It follows that $d = a$, $e = b$ and $f = c$. Hence, this is $D(2, 1, \alpha)^{(1)}$. $\square$

Now that we have found all subfinite regular Kac-Moody extensions of $D(2, 1, \alpha)$ by one vertex, we may restrict our attention to diagrams that do not contain a $D(2, 1, \alpha)$ subdiagram. In particular, the ratio of an isotropic vertex must be a negative rational number by Lemma 2.7, and an isotropic vertex has at most degree two by Corollary 2.8.

Lemma 2.10. *The subfinite regular Kac-Moody 4-vertex extensions of $C(3)$ that are not extensions of $D(2, 1, \alpha)$ are the following: $C(4)$, $C(3)^{(1)}$, $B(1, 2)^{(1)}$, and $A(1, 4)^{(2)}$.*

Proof. Since an isotropic vertex has degree at most two, we are reduced to the following case.




◊ If v_4 is $\otimes$, then $\Gamma_2 \in \mathcal{F}$ implies $b = -1$. This is $D(2, 2)$.

◊ If v_4 is $\bullet$, then $\Gamma_2 \in \mathcal{F}$ implies $a = -2$, $b = -1$. This is $B(1, 2)^{(1)}$.

◊ If v_4 is $\circ$, then $\Gamma_2 \in \mathcal{F}$ implies $a = -2$, $b = -1$. This is $A(1, 4)^{(2)}$.

◊ If v_4 is $\circ$, then $\Gamma_2 \in \mathcal{F}$ implies $a = -1$, $b \in \{-1, -2, -3\}$. If $b = -1$, then this is $C(4)$. If $b = -2$, then this is $C(3)^{(1)}$. If $b = -3$, then by reflecting at v_2 and then at v_1 we obtain




But, $r_1(r_2(\Gamma))_1 \notin \mathcal{F}$. Hence, $b \neq -3$. $\square$

Lemma 2.11. *All subfinite regular Kac-Moody extensions of $G(3)$ are extensions of $D(2, 1, \alpha)$.*

Proof. Since an isotropic vertex has degree at most two, we are reduced to the


following case.



But, $\Gamma_3 \notin \mathcal{F}$. □

Lemma 2.12. *The subfinite regular Kac-Moody extensions of $B(1, 2)$ that are not extensions of $D(2, 1, \alpha)$, $C(3)$ or $G(3)$ are the following: $B(1, 3)$, $B(2, 2)$, $D(2, 2)^{(2)}$, and $A(2, 4)^{(4)}$.*

Proof. Since $\otimes$ has at most degree two and $\bullet$ has at most degree one, we are reduced to the following case.



◊If v_4 is $\otimes$, then $\Gamma_3 \in \mathcal{F}$ implies $b = -1$. This is $B(2, 2)$.


◊If v_4 is $\bullet$, then $\Gamma_3 \in \mathcal{F}$ implies $a = -2$, $b = -1$. This is $D(2, 2)^{(2)}$.

◊If v_4 is $\circ$, then $\Gamma_3 \in \mathcal{F}$ implies $a \in \{-1, -2\}$ and $b \in \{-1, -2, -3\}$. By assumption, Γ_3 is not $C(3)$ or $G(3)$, which implies $b = -1$. If $a = -1$, then this is $B(1, 3)$. If $a = -2$, then this is $A(2, 4)^{(4)}$. □

Lemma 2.13. *The subfinite regular Kac-Moody extensions of $B(2, 1)$ that are not extensions of $D(2, 1, \alpha)$, $C(3)$ or $G(3)$ are the following: $B(3, 1)$, $B(2, 2)$, $A(2, 4)^{(4)}$, and $D(1, 3)^{(2)}$.*


Proof. We consider each case for attaching a vertex to a $B(2, 1)$ diagram.

Case 1:



Then $\Gamma_1 \notin \mathcal{F}$ even if $c, d = 0$.

Case 2:



◊If v_4 is $\otimes$, then $\Gamma_3 \in \mathcal{F}$ implies $b = -1$. This is $B(2, 2)$.

◊If v_4 is $\bullet$, then $\Gamma_3 \in \mathcal{F}$ implies $a = -2$, $b = -1$. This is $A(2, 4)^{(4)}$.

◊If v_4 is $\bigcirc$, then $\Gamma_3 \in \mathcal{F}$ implies $a \in \{-1, -2\}$ and $b \in \{-1, -2, -3\}$. By assumption, Γ_3 is not $C(3)$ or $G(3)$, which implies $b = -1$. If $a = -1$, then this is $B(3, 1)$. If $a = -2$, then this is $D(1, 3)^{(2)}$. $\square$

Lemma 2.14. *The subfinite regular Kac-Moody extensions of $A(0, 2)$ that are not extensions of $D(2, 1, \alpha)$, $C(3)$ or $G(3)$ are the following: $A(0, 3)$, $A(1, 2)$, $B(1, 3)$, $B(3, 1)$, $A(0, 2)^{(1)}$, and $q(4)^{(2)}$.*

Proof. We consider each case for attaching a vertex to an $A(0, 2)$ diagram.

Case 1:

$$\bigvee_4 \xrightleftharpoons[b]{a} \bigotimes_1 \xrightleftharpoons[1]{1} \bigotimes_2 \xrightleftharpoons[-1]{-1} \bigcirc_3 \quad a, b \neq 0$$

◊If v_4 is $\bigotimes$, then $\Gamma_3 \in \mathcal{F}$ implies $b = -1$. This is $A(1, 2)$.

◊If v_4 is $\bullet$, then $\Gamma_3 \in \mathcal{F}$ implies $a = -2$, $b = -1$. This is $B(1, 3)$.

◊If v_4 is $\bigcirc$, then $\Gamma_3 \in \mathcal{F}$ implies $a \in \{-1, -2\}$ and $b \in \{-1, -2, -3\}$. By assumption, Γ_3 is not $C(3)$ or $G(3)$, which implies $b = -1$. If $a = -1$, then this is $A(0, 3)$. If $a = -2$, then this is $B(3, 1)$.

Case 2:

$$\bigotimes_1 \xrightleftharpoons[1]{1} \bigotimes_2 \xrightleftharpoons[-1]{-1} \bigcirc_3 \xrightleftharpoons[a]{b} \bigvee_4 \quad a, b \neq 0$$

◊If v_4 is $\bigotimes$, then $\Gamma_1 \in \mathcal{F}$ implies $b = -1$. This is $A(1, 2)$.

◊If v_4 is $\bullet$, then $\Gamma_1 \in \mathcal{F}$ implies $a = -2$, $b = -1$. This is $B(1, 3)$.

◊If v_4 is $\bigcirc$, then $\Gamma_1 \in \mathcal{F}$ implies $a \in \{-1, -2\}$ and $b \in \{-1, -2, -3\}$. By assumption, Γ_1 is not $C(3)$ or $G(3)$, which implies $b = -1$. If $a = -1$, then this is $A(0, 3)$. If $a = -2$, then this is $B(3, 1)$.

Case 3:

$$\begin{array}{ccc} \bigotimes_2 & \xrightleftharpoons[-1]{-1} & \bigcirc_3 \\ \uparrow 1 & & \uparrow b \\ \bigotimes_1 & \xrightleftharpoons[c]{d} & \bigvee_4 \end{array} \quad a, b, c, d \neq 0$$

◊If v_4 is $\bigcirc$, then $\Gamma_2 \in \mathcal{F}$ implies $c = -1$ and $b \in \{-1, -2, -3\}$. By assumption, Γ_2 is not $C(3)$ or $G(3)$, which implies $b = -1$. Since we assume that Γ_3 is not $C(3)$ or $G(3)$, we have that $\Gamma_3 \in \mathcal{F}$ implies $d = -1$. Now $\Gamma_1 \in \mathcal{F}$ implies $a \in \{-1, -2, -3\}$ and since we assume that it is not $C(3)$ or $G(3)$, we have $a = -1$. This is $A(0, 2)^{(1)}$.

◊If v_4 is $\bigotimes$, then $\Gamma_1 \in \mathcal{F}$ implies $a = -1$. Since we assume that Γ_3 is not $D(2, 1, \alpha)$, we have that $\Gamma_3 \in \mathcal{F}$ implies $d = -1$. Finally, $\Gamma_2 \in \mathcal{F}$ and not equal to $C(3)$ or $G(3)$ implies $\frac{b}{c} = -1$. This is $q(4)^{(2)}$. $\square$

Lemma 2.15. *The subfinite regular Kac-Moody extensions of $A(1,1)$ that are not extensions of $D(2,1,\alpha)$, $C(3)$ or $G(3)$ are the following: $A(1,2)$, $B(2,2)$, $A(1,1)^{(1)}$, and $q(4)^{(2)}$.*

Proof. We consider each case for attaching a vertex to an $A(1,1)$ diagram.

Case 1:

$$\otimes_1 \xrightarrow[1]{1} \otimes_2 \xleftarrow[1]{-1} \otimes_3 \xrightleftharpoons[a]{b} \odot_4 \quad a, b \neq 0$$

◊If v_4 is $\otimes$, then $\Gamma_1 \in \mathcal{F}$ implies $b = -1$. This is $A(1,2)$.

◊If v_4 is $\bullet$, then $\Gamma_1 \in \mathcal{F}$ implies $a = -2$, $b = -1$. This is $B(2,2)$.

◊If v_4 is $\circ$, then $\Gamma_1 \in \mathcal{F}$ implies $a \in \{-1, -2\}$ and $b \in \{-1, -2, -3\}$. By assumption, Γ_1 is not $C(3)$ or $G(3)$, which implies $b = -1$. If $a = -1$, then this is $A(1,2)$. If $a = -2$, then this is $B(2,2)$.

Case 2:

$$\begin{array}{ccc} \otimes_1 & \xrightleftharpoons[1]{-1} & \otimes_2 \\ \uparrow 1 & & \uparrow b \\ \otimes_3 & \xrightleftharpoons[c]{d} & \odot_4 \end{array} \quad a, b, c, d \neq 0$$

◊If v_4 is $\otimes$, then $\Gamma_1 \in \mathcal{F}$ implies $a = -1$, $\Gamma_3 \in \mathcal{F}$ implies $d = -1$, and $\Gamma_2 \in \mathcal{F}$ implies $\frac{b}{c} = -1$. This is $A(1,1)^{(1)}$.

◊If v_4 is $\circ$, then $\Gamma_1 \in \mathcal{F}$ implies $b = c = -1$. Then $\Gamma_2, \Gamma_3 \in \mathcal{F}$ and not $C(3)$ or $G(3)$ imply $a = d = -1$. This is $q(4)^{(2)}$. $\square$

This completes the classification of connected subfinite regular Kac-Moody diagrams with four vertices. We observe that all are either of finite type or have finite growth.

2.4 Subfinite regular Kac-Moody: 5 vertices

For each finite type 4-vertex diagram, we will consider each case for attaching a vertex to the diagram. Let v_5 denote the additional vertex, and let Γ denote the corresponding extended diagram. Recall that a_{ij} denotes the label of the arrow from the vertex v_i to the vertex v_j . Also, $a_{ij} = 0$ if and only if $a_{ji} = 0$.

Lemma 2.16. *The subfinite regular Kac-Moody extensions of $D(2,2)$ are the following: $D(2,3)$, $D(3,2)$, $B(2,2)^{(1)}$, $A(4,3)^{(2)}$, $A(3,3)^{(2)}$, $D(2,2)^{(2)}$.*

Proof.

$$\begin{array}{ccccc} & & \otimes_2 & & \\ & \nearrow 1 & & \nwarrow -2 & \\ \otimes_4 & \xleftarrow[-1]{-1} & \otimes_1 & \xrightleftharpoons[1]{1} & \otimes_3 \end{array}$$

First observe that $\Gamma_4 \in \mathcal{F}$ implies $a_{52} = a_{53} = 0$.


◊If v_5 is $\bullet$, then $\Gamma_2 \in \mathcal{F}$ implies $a_{51} = 0$, $a_{54} = -2$, $a_{45} = -1$. This is $B(2, 2)^{(1)}$.

◊If v_5 is $\circ$, then $\Gamma_2 \in \mathcal{F}$ implies $a_{51} = 0$ and $a_{45}, a_{54} \in \{-1, -2\}$. If $a_{54} = -2$ and $a_{45} = -1$, then this is $A(4, 3)^{(2)}$. If $a_{54} = -1$ and $a_{45} = -2$, then this is $D(2, 2)^{(2)}$. If $a_{54} = a_{45} = -1$, then this is $D(2, 3)$.

◊If v_5 is $\otimes$, then $\Gamma_2 \in \mathcal{F}$ implies that either $a_{51} = a_{15} = 0$, $a_{45} = -1$, $a_{54} = 1$ and this is $D(3, 2)$, or $a_{51} = -1$, $a_{15} = 1$, $a_{45} = a_{54} = -2$ and this is $A(3, 3)^{(2)}$. $\square$

Lemma 2.17. *The subfinite regular Kac-Moody extensions of $D(3, 1)$ are the following: $D(2, 3)$, $D(4, 1)$, $B(3, 1)^{(1)}$, $A(2, 5)^{(2)}$, $A(3, 3)^{(2)}$, $A(5, 1)^{(2)}$.*

Proof.



First observe that $\Gamma_4 \in \mathcal{F}$ implies $a_{52} = a_{53} = 0$.


◊If v_5 is $\bullet$, then $\Gamma_3 \in \mathcal{F}$ implies $a_{51} = 0$, $a_{54} = -2$, $a_{45} = -1$. This is $A(2, 5)^{(2)}$.

◊If v_5 is $\circ$, then $\Gamma_3 \in \mathcal{F}$ implies that Γ satisfies the following conditions. If $a_{51} \neq 0$, then $a_{51} = a_{15} = -1$, $a_{54} = a_{45} = 0$, and this is $A(3, 3)^{(2)}$. If $a_{51} = 0$, then $a_{54}, a_{45} \in \{-1, -2\}$. If $a_{51} = 0$, $a_{54} = -2$ and $a_{45} = -1$, then this is $B(3, 1)^{(1)}$. If $a_{51} = 0$, $a_{54} = -1$ and $a_{45} = -2$, then this is $A(5, 1)^{(2)}$. If $a_{51} = 0$, $a_{54} = -1$ and $a_{45} = -1$, then this is $D(4, 1)$.

◊If v_5 is $\otimes$, then $\Gamma_3 \in \mathcal{F}$ implies $a_{51} = 0$, $a_{45} = -1$, $a_{54} = 1$. This is $D(2, 3)$. $\square$

Lemma 2.18. *The subfinite regular Kac-Moody extensions of $C(4)$ are the following: $C(5)$, $D(2, 3)$, $B(1, 3)^{(1)}$, $D(1, 3)^{(1)}$, $A(5, 1)^{(2)}$, $A(6, 1)^{(2)}$.*

Proof.



First observe that $\Gamma_4 \in \mathcal{F}$ implies $a_{52} = a_{53} = 0$.


◊If v_5 is $\bullet$, then $\Gamma_3 \in \mathcal{F}$ implies $a_{51} = 0$, $a_{54} = -2$, $a_{45} = -1$. This is $B(1, 3)^{(1)}$.

◊If v_5 is $\circ$, then $\Gamma_3 \in \mathcal{F}$ implies that Γ satisfies the following conditions. If $a_{51} \neq 0$, then $a_{51} = a_{15} = -1$, $a_{54} = a_{45} = 0$, and this is $A(5, 1)^{(2)}$. If $a_{51} = 0$,

then $a_{54}, a_{45} \in \{-1, -2\}$. If $a_{51} = 0$, $a_{54} = -2$ and $a_{45} = -1$, then this is $A(6, 1)^{(2)}$. If $a_{51} = a_{15} = 0$, $a_{54} = -1$ and $a_{45} = -2$, then this is $D(1, 3)^{(1)}$. If $a_{51} = a_{15} = 0$, $a_{54} = -1$ and $a_{45} = -1$, then this is $C(5)$.
 $\diamond$ If v_5 is $\otimes$, then $\Gamma_3 \in \mathcal{F}$ implies $a_{51} = 0$, $a_{54} = 1$, $a_{45} = -1$. This is $D(2, 3)$. $\square$

Lemma 2.19. *The only subfinite regular Kac-Moody extensions of $F(4)$ is $F(4)^{(1)}$.*

Proof.



First observe that $\Gamma_4 \in \mathcal{F}$ implies $a_{52} = a_{53} = 0$. Then $\Gamma_2 \in \mathcal{F}$ implies $a_{51} = 0$, $a_{54} = a_{45} = -1$. This is $F(4)^{(1)}$. $\square$

Let $\mathcal{K} = \{\Gamma' \in \mathcal{F} \mid R(\Gamma') \text{ is a chain for every sequence } R \text{ of odd reflections}\}$.

Lemma 2.20. *The subfinite regular Kac-Moody extensions of $A(0, 3)$, $A(1, 2)$, $B(1, 3)$, $B(3, 1)$ and $B(2, 2)$, that are not extensions of $C(4)$, $D(3, 1)$, $D(2, 2)$ or F_4 , are the following: $A(0, 4)$, $A(1, 3)$, $A(2, 2)$, $B(1, 4)$, $B(2, 3)$, $B(3, 2)$, $B(4, 1)$, $A(0, 3)^{(1)}$, $A(1, 2)^{(1)}$, $A(4, 4)^{(2)}$, $A(6, 2)^{(2)}$, $D(2, 3)^{(2)}$, $D(3, 2)^{(2)}$, $D(4, 1)^{(2)}$, $q(5)^{(2)}$.*

Proof. Note that every connected 4-vertex subdiagram of Γ containing an isotropic vertex is an element of $\mathcal{K}$. We find the extensions for each of the diagrams: $A(0, 3)$, $A(1, 2)$, $B(1, 3)$, $B(3, 1)$ and $B(2, 2)$.

$$B(3, 1) \quad \otimes_1 \xrightleftharpoons[1]{1} \otimes_2 \xrightleftharpoons[-1]{-1} \bigcirc_3 \xrightleftharpoons[-2]{-1} \bigcirc_4$$

First observe that since $\Gamma_1, \Gamma_4 \in \mathcal{K}$ we have $a_{52} = a_{53} = a_{54} = 0$.

$\diamond$ If v_5 is $\otimes$, then $\Gamma_4 \in \mathcal{K}$ implies $a_{15} = -1$, $a_{15} = 1$. This is $B(3, 2)$.

$\diamond$ If v_5 is $\bullet$, then $\Gamma_4 \in \mathcal{K}$ implies $a_{15} = -1$, $a_{51} = -2$. This is $A(6, 2)^{(2)}$.

$\diamond$ If v_5 is $\bigcirc$, then $\Gamma_4 \in \mathcal{K}$ implies $a_{15} = -1$, $a_{51} \in \{-1, -2\}$. If $a_{51} = -1$, then this is $B(4, 1)$. If $a_{51} = -2$, then this is $D(4, 1)^{(2)}$.

$$B(1, 3) \quad \otimes_1 \xrightleftharpoons[1]{1} \otimes_2 \xrightleftharpoons[-1]{-1} \bigcirc_3 \xrightleftharpoons[-2]{-1} \bullet_4$$

First observe that since $\Gamma_1, \Gamma_4 \in \mathcal{K}$ we have $a_{52} = a_{53} = a_{54} = 0$.

$\diamond$ If v_5 is $\otimes$, then $\Gamma_4 \in \mathcal{K}$ implies $a_{15} = -1$, $a_{15} = 1$. This is $B(2, 3)$.

$\diamond$ If v_5 is $\bullet$, then $\Gamma_4 \in \mathcal{K}$ implies $a_{15} = -1$, $a_{51} = -2$. This is $D(2, 3)^{(2)}$.

$\diamond$ If v_5 is $\bigcirc$, then $\Gamma_4 \in \mathcal{K}$ implies $a_{15} = -1$, $a_{51} \in \{-1, -2\}$. If $a_{51} = -1$, then

this is $B(1, 4)$. If $a_{51} = -2$, then this is $A(6, 2)^{(2)}$.

$$B(2, 2) \quad \otimes_1 \xrightleftharpoons[1]{1} \otimes_2 \xrightleftharpoons[1]{-1} \otimes_3 \xrightleftharpoons[-2]{-1} \bullet_4$$

First observe that since $\Gamma_1, \Gamma_4 \in \mathcal{K}$ we have $a_{52} = a_{53} = a_{54} = 0$.

◊ If v_5 is $\otimes$, then $\Gamma_4 \in \mathcal{K}$ implies $a_{15} = -1, a_{15} = 1$. This is $B(2, 3)$.

◊ If v_5 is $\bullet$, then $\Gamma_4 \in \mathcal{K}$ implies $a_{15} = -1, a_{51} = -2$. This is $A(4, 4)^{(2)}$.

◊ If v_5 is $\circ$, then $\Gamma_4 \in \mathcal{K}$ implies $a_{15} = -1, a_{51} \in \{-1, -2\}$. If $a_{51} = -1$, then this is $B(3, 2)$. If $a_{51} = -2$, then this is $D(3, 2)^{(2)}$.

Finally, we restrict our attention to diagrams satisfying: a connected 4-vertex subdiagram containing an isotropic vertex is a diagram for $A(0, 3)$ or $A(1, 2)$.

$$A(1, 2) \quad \otimes_1 \xrightleftharpoons[1]{1} \otimes_2 \xrightleftharpoons[1]{-1} \otimes_3 \xrightleftharpoons[1]{-1} \otimes_4$$

First observe that since $\Gamma_1, \Gamma_4 \in \mathcal{K}$ we have $a_{52} = a_{53} = 0$. There are two distinct cases: $a_{54} = 0$ and $a_{51}, a_{54} \neq 0$.

◊ If v_5 is $\otimes$ and $a_{54} = 0$, then $a_{51} = 1, a_{15} = -1$. This is $A(2, 2)$.

◊ If v_5 is $\circ$ and $a_{54} = 0$, then $a_{51}, a_{15} = -1$. This is $A(1, 3)$.

◊ If v_5 is $\otimes$ and $a_{54}, a_{51} \neq 0$, then $a_{51} = 1, a_{15}, a_{54}, a_{45} = -1$. This is $q(5)^{(2)}$.

◊ If v_5 is $\circ$ and $a_{54}, a_{51} \neq 0$, then $a_{51}, a_{15}, a_{54}, a_{45} = -1$. This is $A(1, 2)^{(1)}$.

$$A(0, 3) \quad \circ_1 \xrightleftharpoons[1]{-1} \otimes_2 \xrightleftharpoons[1]{-1} \otimes_3 \xrightleftharpoons[-1]{-1} \circ_4$$

Now we assume that a connected 4-vertex subdiagram is a diagram for $A(0, 3)$.

First observe that since $\Gamma_1, \Gamma_4 \in \mathcal{K}$ we have $a_{52}, a_{53} = 0$. There are two distinct cases: $a_{54} = 0$ and $a_{51}, a_{54} \neq 0$.

◊ If v_5 is $\otimes$ and $a_{54} = 0$, then $a_{51} = 1, a_{15} = -1$. This is $A(1, 3)$.

◊ If v_5 is $\circ$ and $a_{54} = 0$, then $a_{51}, a_{15} = -1$. This is $A(0, 4)$.

◊ If v_5 is $\otimes$ and $a_{54}, a_{51} \neq 0$, then $a_{51} = 1, a_{15}, a_{54}, a_{45} = -1$. This is $q(5)^{(2)}$.

◊ If v_5 is $\circ$ and $a_{54}, a_{51} \neq 0$, then $a_{51}, a_{15}, a_{54}, a_{45} = -1$. This is $A(0, 3)^{(1)}$. $\square$

This completes the classification of connected subfinite regular Kac-Moody diagrams with five vertices. We observe that all are either of finite type or have finite growth.

2.5 Subfinite regular Kac-Moody: $n \geq 6$ vertices

Now we handle the general case. We find all connected subfinite regular Kac-Moody diagrams with six or more vertices. We will show in Theorem 5.5 that the subfinite regular Kac-Moody diagrams which are not of finite type are not extendable, completing the classification of regular Kac-Moody diagrams.

Remark 2.21. A finite type diagram Γ' with $n \geq 5$ vertices has the following form:

$$\begin{array}{c} \bullet 2 \\ \swarrow \quad \searrow \\ \circledast 1 \quad \bullet 3 \text{ --- } \bullet_{n-2} \text{ --- } \bullet_{n-1} \end{array}, \quad (8)$$

where Γ'_1 is $A(k, l)$ or A_k , and the subdiagram $\{v_1, v_2, v_3\}$ is one of the diagrams in the following table.

Table 2.21

Lemma 2.22. Let Γ be a subfinite regular Kac-Moody cycle with $n \geq 4$ vertices and let v_j be a vertex of Γ connected to v_i and v_k . Then either

$$\begin{array}{l} v_j \text{ is } \circ \text{ with } a_{ji} = a_{jk} = -1; \text{ or} \\ v_j \text{ is } \otimes \text{ with } \frac{a_{ji}}{a_{jk}} = -1. \end{array} \quad (9)$$

If Γ has an even number of odd roots, then $\mathfrak{g}(A)$ is an $A(k-1, l)^{(1)}$ diagram. If Γ has an odd number of odd roots, then $\mathfrak{g}(A)$ is a $q(n)^{(2)}$ diagram.


Proof. By the above classification, all subfinite regular Kac-Moody 4-vertex and 5-vertex cycles are of type $A(k, l)^{(1)}$ or $q(n)^{(2)}$, and satisfy (9). Let Γ be a subfinite regular Kac-Moody cycle with $n \geq 6$ vertices and let v_j be a vertex which is connected to v_i and v_k . Since $n \geq 6$, v_j is contained in a proper 5-vertex subdiagram where v_j is the middle vertex of the chain. By the 5-vertex classification, we see that for every finite type chain diagram with five vertices, the middle vertex satisfies (9). Hence, every vertex of Γ satisfies (9). If Γ has an even number of odd roots, then the corresponding matrix A is symmetrizable

and $\mathfrak{g}(A)$ is an $A(k, l)^{(1)}$ diagram. If Γ has an odd number of odd roots, then A is non-symmetrizable and $\mathfrak{g}(A)$ is a $q(n)^{(2)}$ diagram. $\square$

Lemma 2.23. *If Γ contains a proper subdiagram Γ' which is a cycle with $n \geq 4$ vertices then Γ is not subfinite regular Kac-Moody.*


Proof. Suppose Γ is a subfinite regular Kac-Moody diagram which contains a cycle with $n \geq 4$ vertices. By the Lemma 2.5, we are reduced to the case where $\Gamma = \Gamma' \cup \{v_{n+1}\}$ is subfinite regular Kac-Moody. By the above classification, a subfinite regular Kac-Moody 5-vertex diagram does not contain a 4-vertex cycle subdiagram. So now suppose Γ' is a cycle with n vertices where $n \geq 5$ and that $\Gamma = \Gamma' \cup \{v_{n+1}\}$ is subfinite regular Kac-Moody. If Γ' contains an isotropic vertex, then by Lemma 2.22 the diagram Γ' is either $A(m, n)^{(1)}$ or $q(n)^{(2)}$, which is not of finite type.

Suppose Γ' does not contain an isotropic vertex. Then the additional vertex v_{n+1} of Γ is isotropic. Since Γ is connected we have the following subdiagram:



where the double lines are necessarily connected, and the dotted lines are possibly connected. But by the 5-vertex classification, we see that this is not a finite type subdiagram. $\square$


Lemma 2.24. *Let Γ be the following diagram*



where Γ contains an isotropic vertex and Γ_6 is a diagram for $A(k, l)$ or A_k . Then Γ is not a subfinite regular Kac-Moody diagram.

Proof. If v_6 is isotropic, then $\Gamma_5 \notin \mathcal{F}$, which is a contradiction. We may assume that v_3 is isotropic, by using odd reflections in the subdiagram $\Gamma_{3,6}$. Then $\Gamma_1, \Gamma_5 \in \mathcal{F}$ implies that the vertices v_2, v_4, v_6 are even. Moreover, $\Gamma_1 \in \mathcal{F}$ implies that the ratio of v_3 in the subdiagram $\{v_2, v_3, v_6\}$ is 1, while $\Gamma_5 \in \mathcal{F}$ implies that the ratio is -1 , which is a contradiction. $\square$

Proposition 2.25. *If Γ is a subfinite regular Kac-Moody diagram with $n \geq 6$ vertices and Γ is not a cycle, then Γ is of the form*



where the subdiagram $\Gamma \setminus \{v_1, v_n\}$ is $A(k, l)$ or A_k , and the subdiagrams $\{v_1, v_2, v_3\}$ and $\{v_n, v_{n-1}, v_{n-2}\}$ are diagrams from Table 2.21.

Proof. First note that diagrams satisfying (10) reflect by odd reflections to diagrams that again satisfy (10). So it suffices to prove the proposition for a diagram obtained by odd reflections from the original diagram.

Let Γ be a subfinite regular Kac-Moody diagram with $n \geq 6$ vertices, which is not a cycle. Choose a vertex v_n such that $\Gamma' := \Gamma \setminus \{v_n\}$ is connected and contains an isotropic vertex. Then Γ' satisfies (8). We may assume that the subdiagram $\{v_1, v_2, v_3\}$ contains an isotropic vertex, by using odd reflections in the subdiagram $\{v_4, \dots, v_{n-1}\}$. If $\Gamma' \setminus \{v_1\}$ does not contain an isotropic vertex, then v_1 is isotropic. In this case, by Table 2.21, Γ' is the standard diagram for $A(1, n-2)$ and an odd reflection at v_1 results in a diagram with v_2 isotropic. So we may assume that v_2 or v_3 is isotropic.

Suppose that v_n is connected only to the vertex v_1 . If $a_{12}, a_{21} = 0$, then $a_{13}, a_{31} \neq 0$ and $\{v_n, v_1, v_3, v_2, v_4\} \in \mathcal{F}$ implies that the subdiagram $\{v_n, v_1, v_3\}$ is $A(k, l)$ or A_k . Then by Lemma 2.24, the diagram $\{v_n, v_1, v_2, v_3, v_4, v_5\}$ is not a subfinite regular Kac-Moody diagram. If $a_{12}, a_{21} \neq 0$, then $\Gamma_{n-1} \in \mathcal{F}$ implies $a_{13}, a_{31} = 0$. The condition $\{v_n, v_1, v_2, v_3, v_4\} \in \mathcal{F}$ yields two possibilities: either $\{v_n, v_1, v_2\}$ is a diagram from Table 2.21 and $\{v_1, v_2, v_3, v_4\}$ is $A(k, l)$ or A_k , or $\{v_4, v_3, v_2\}$ is a diagram from Table 2.21 and $\{v_n, v_1, v_2, v_3\}$ is $A(k, l)$ or A_k . In the first case, Γ_n is $A(k, l)$ or A_k , and we are done. In the second case, Γ is $A(k, l)$ or A_k , and we are done.

Now we may assume that $\Gamma \setminus \{v_1\}$ is connected, and hence satisfies (8). Thus, the vertex v_n is not connected to vertices v_j with $3 < j < n-2$. First suppose that v_n is connected to v_{n-2} or v_{n-1} . Then $\Gamma_1 \in \mathcal{F}$ implies that v_n is not connected to v_2 or v_3 and Γ_1 satisfies (8). By Lemma 2.23, v_n is not connected to v_1 . Thus, Γ satisfies (10). Now suppose that v_n is not connected to v_{n-2} or v_{n-1} . If v_n is connected to v_3 then $\{v_n, v_1, v_2, v_3, v_4\}$ satisfies (8), implying v_n is not connected to v_1 or v_2 , $a_{13}, a_{31} = 0$, and the subdiagram $\{v_1, v_2, v_3\}$ is $A(k, l)$ or A_k . But then by Lemma 2.24, $\{v_n, v_1, v_2, v_3, v_4, v_5\}$ is not a subfinite regular Kac-Moody diagram. Hence, v_n is not connected to v_3 . Finally, since subdiagram $\Gamma \setminus \{v_{n-1}\}$ satisfies (8) with v_n connected only to v_1 or v_2 we conclude that Γ satisfies (10). $\square$


2.6 Classification theorem

Theorem 2.26. *If A is a symmetrizable matrix and $\mathfrak{g}(A)$ has a simple isotropic root, then $\mathfrak{g}(A)$ is regular Kac-Moody if and only if it has finite growth.*

If A is a non-symmetrizable matrix and $\mathfrak{g}(A)$ has a simple isotropic root, then $\mathfrak{g}(A)$ is regular Kac-Moody if and only if it is one of the following three classes:

Algebra	Dynkin diagrams
$q(n)^{(2)}$	There are n $\bullet$. Each $\bullet$ is either $\bigcirc$ or $\otimes$. An odd number of them are $\otimes$. If $\bullet$ is $\bigcirc$, then $a = b = -1$. If $\bullet$ is $\otimes$, then $\frac{a}{b} = -1$.
$S(1, 2, \alpha)$	$\alpha \neq 0, \frac{1}{\alpha} \in \mathbb{C} \setminus \mathbb{Z}$
$Q^\pm(m, n, t)$	$1 + a + \frac{1}{b} = m$ $1 + b + \frac{1}{c} = n$ $1 + c + \frac{1}{a} = t$ $m, n, t \in \mathbb{Z}_{\leq -1}$ and not all equal to -1, $a, b, c \in \mathbb{R} \setminus \mathbb{Q}$.

3 The Lie superalgebra $Q^\pm(m, n, t)$



In this section, we describe the parameters of the defining matrices for $Q^\pm(m, n, t)$, and then we show that $Q^\pm(m, n, t)$ is not symmetrizable and does not have finite growth.

Lemma 3.1. *For each diagram $Q^\pm(m, n, t)$, it follows that $a, b, c \in \mathbb{R} \setminus \mathbb{Q}$, and there are two solutions of the above equations, namely $Q^-(m, n, t)$ with $a, b, c < -1$ and $Q^+(m, n, t)$ with $-1 < a, b, c < 0$.*

Proof. First, we show that a solution exists. To simplify the calculations we let

$$M = 1 - m, \quad N = 1 - n, \quad T = 1 - t.$$

Then $M, N, T \in \mathbb{Z}_{\geq 2}$. Solving equation 1 for b and equation 3 for c , and then substituting into equation 2 yields

$$-N = \frac{-1}{a + M} + \frac{-a}{aT + 1}.$$

By clearing denominators and regrouping, we have $f(a) = 0$ where

$$f(a) = (NT - 1)a^2 + (MNT - M + N - T)a + (MN - 1).$$

Now $(NT - 1) \neq 0$, because $N, T \geq 2$. So this is a quadratic equation in the variable a with integer coefficients. We solve the equation $f(a) = 0$ for a . By applying the quadratic formula, we have

$$a = \frac{-B \pm \sqrt{B^2 - 4AC}}{2A}, \quad A = NT - 1, \quad B = MNT - M + N - T, \quad C = MN - 1.$$

Next we show that the discriminant is strictly positive, implying that there are two real solutions. Now

$$B^2 - 4AC = (MNT - M + N - T)^2 - 4(NT - 1)(MN - 1) = (MNT - M - N - T)^2 - 4.$$

So we only need to show that $(MNT - M - N - T) > 2$, which follows from the following calculation. Since $M - 1, N - 1, T - 1 \geq 1$ and not all equal to 1, we have

$$\begin{aligned} 1 &< (M - 1)(N - 1)(T - 1) \\ &= MNT - (N - 1)M - (T - 1)N - (M - 1)T - 1 \\ &< MNT - M - N - T - 1. \end{aligned}$$

Hence, $(MNT - M - N - T) > 2$ and the discriminant is strictly positive. Thus, $a \in \mathbb{Q}(\sqrt{B^2 - 4AC}) \subset \mathbb{R}$. By the defining equations, it follows that $a, b, c \in \mathbb{Q}(\sqrt{B^2 - 4AC}) \subset \mathbb{R}$.

Next we show that $a, b, c \leq 0$, and either $-1 < a, b, c < 0$ or $a, b, c \leq -1$. First note that $a, b, c \neq 0$. Now if $a, b > 0$ then $-M = a + \frac{1}{b} > 0$, but $-M < 0$. So without loss of generality by symmetry, we have $a, b < 0$. If $a \leq -1$, then $0 < \frac{-1}{a} \leq 1$ which implies $c = \frac{-1}{a} - T \leq 1 - T \leq -1$. If $a > -1$, then $a + M > M - 1 \geq 1$. This implies $b = \frac{-1}{a+M} > -1$ and

$$N(a + M) - 1 \geq 2(a + M) - 1 = (a + M) + ((a + M) - 1) \geq (a + M) > 1,$$

which implies $-1 < c < 0$ since

$$c = \frac{-1}{b + N} = \frac{-1}{\frac{-1}{a+M} + N} = \frac{-(a + M)}{N(a + M) - 1}.$$

So in both cases $c < 0$. If $-1 < a < 0$, then $-1 < b, c < 0$. By the symmetry of the equations, we conclude that either $-1 < a, b, c < 0$ or $a, b, c \leq -1$. Now if $a = -1$ then $b, c \leq -1$. From the first equation we obtain $\frac{1}{b} = m = -1$. Then from the second equation, $\frac{1}{c} = n = -1$. But then $a, b, c = -1$ implies $m, n, t = -1$ and this diagram defines $q(3)^{(2)}$, not $Q^\pm(m, n, t)$. Therefore, either $-1 < a, b, c < 0$ or $a, b, c < -1$.

Next we show that a, b, c are not rational. It is easy to see from the equations that if one of them is rational then they all are. Now suppose we have a rational solution

$$a = -\frac{r}{s}, b = -\frac{t}{u}, c = -\frac{v}{w}$$

where $r, s, t, u, v, w \in \mathbb{Z}_{>0}$ and each fraction is reduced, ie. $\gcd(r, s) = 1$, $\gcd(t, u) = 1$, $\gcd(v, w) = 1$. Then

$$-M = a + \frac{1}{b} = -\frac{r}{s} - \frac{u}{t} = -\frac{rt + us}{st}.$$

So, $stM = rt + us$. Then $s(Mt - u) = rt$, implying s divides rt . Since $\gcd(r, s) = 1$, we have that s divides t . Also, $t(Ms - r) = us$ implies t divides us . Since $\gcd(t, u) = 1$, we have that t divides s . Hence, $s = t$. Similarly, by the same type of calculation we conclude that $u = v$ and $w = r$. Making these substitutions into the above equations we have

$$a = -\frac{r}{s}, b = -\frac{s}{v}, c = -\frac{v}{r}.$$

We see that if $-1 < a, b, c < 0$ then $r < s$, $s < v$, and $v < r$, which is a contradiction. If $a, b, c < -1$, then $r > s$, $s > v$, and $v > r$, which is again a contradiction. Hence, there is no rational solution. Therefore, $a, b, c \in \mathbb{R} \setminus \mathbb{Q}$.

For each $m, n, t \in \mathbb{Z}_{\leq -1}$ and not all equal to -1 , we have shown that we have two real non-rational solutions. Now

$$f(a) = (NT - 1)a^2 + (MNT - M + N - T)a + (MN - 1),$$

and

$$\begin{aligned}
f(-1) &= (NT - 1) - (MNT + M - N + T) + (MN - 1) \\
&= (1 - M)(N(T - 1) - 1) + (T - 1) \\
&\leq (-1)(N(T - 1) - 1) + (T - 1) \\
&= (1 - N)(T - 1) + 1 \\
&\leq (-1)(T - 1) + 1 \\
&= 2 - T \\
&\leq 0.
\end{aligned}$$

Since -1 is not a solution of $f(a)$ for $Q^\pm(m, n, t)$, we conclude that $f(-1) < 0$. Since the leading coefficient of $f(a)$ is positive, i.e. $NT - 1 > 0$, the quadratic equation $f(a)$ is concave up. Since $f(0) = MN - 1 > 0$, it follows that both zeros of the equation are negative. Because of the sign change, there is a solution between -1 and 0 . Because the graph is concave up with $f(-1) < f(0)$, the second solution is strictly less than -1 . Hence, for each diagram $Q^\pm(m, n, t)$, we have that $a, b, c \in \mathbb{R} \setminus \mathbb{Q}$, and there are two real non-rational solutions of the defining matrix equations, namely $Q^-(m, n, t)$ with $a, b, c < -1$ and $Q^+(m, n, t)$ with $-1 < a, b, c < 0$. $\square$

Corollary 3.2. *The determinant of the Cartan matrix equals $1 + abc$ and is nonzero. Hence, the dimension of the Cartan subalgebra is 3.*

It is clear that $Q^\pm(m, n, t) \cong Q^\pm(n, t, m)$ by cyclic permutation of the variables a, b, c . We also have $Q^\pm(m, n, t) \cong Q^\mp(m, t, n)$ by transforming the equations: $a \rightarrow \frac{1}{b}$, $b \rightarrow \frac{1}{a}$, $c \rightarrow \frac{1}{c}$.

Lemma 3.3. *$Q^\pm(m, n, t)$ is not symmetrizable.*

Proof. Suppose that we have a symmetrizable solution a, b, c . We show that this implies $a \in \mathbb{Q}$, which is a contradiction. If the matrix is symmetrizable then $abc = 1$. So, we substitute $c = \frac{1}{ab}$ into the defining equations. This yields

$$\begin{aligned}
1 + a + \frac{1}{b} &= m \\
1 + (a + 1)b &= n \\
1 + \frac{1}{ab} + \frac{1}{a} &= t
\end{aligned}$$

From the first equation we have $b = \frac{1}{m-a-1}$. Substituting this into the second equation and solving for a , we have $a = \frac{mn-m-n}{n} \in \mathbb{Q}$. Hence, there is no symmetrizable solution. $\square$

Lemma 3.4. $Q^\pm(m, n, t)$ does not have finite growth.

Proof. The set of principal roots of $Q^\pm(m, n, t)$ is $\Pi_0 = \{\alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_3 + \alpha_1\}$. The Cartan matrix B of the subalgebra of $Q^\pm(m, n, t)$ generated by Π_0 is

$$B = \begin{pmatrix} 2 & m & m \\ n & 2 & n \\ t & t & 2 \end{pmatrix}.$$

All off diagonal entries of B are negative integers. Since they are not all equal to -1 , this is not a Cartan matrix of a finite growth Kac-Moody algebra. By Lemma 1.8, $Q^\pm(m, n, t)$ does not have finite growth. $\square$

A Kac-Moody superalgebra is called *hyperbolic* if after the removal of any simple root the superalgebra is either of finite or affine type.

Lemma 3.5. The regular Kac-Moody superalgebra $Q^\pm(m, n, t)$ is hyperbolic for the following $m \geq n \geq t$.

m	n	t
-1	-1	-2
-1	-1	-3
-1	-1	-4
-1	-2	-2
-2	-2	-2

4 Integrable modules

4.1 Integrable modules over affine Lie superalgebras

Let $\mathfrak{g}(A)$ be a regular Kac-Moody superalgebra, and let $\mathfrak{g}(A) = \mathfrak{h} \oplus (\oplus_{\alpha \in \Delta} \mathfrak{g}(A)_\alpha)$ be the root space decomposition. A root α is called *real* if α or $\frac{1}{2}\alpha$ is simple in some reflected base obtained by even and odd reflections, and it is called *imaginary* otherwise. An element $\rho \in \mathfrak{h}^*$ such that $\rho(h_i) = \alpha_i(h_i) = \frac{1}{2}a_{ii}$ for all $i \in I$ is called a *Weyl vector*.

Let $U(\mathfrak{g})$ denote the universal enveloping algebra of a Lie superalgebra $\mathfrak{g}$. Let $\mathfrak{n}_+$ (resp. $\mathfrak{n}_-$) denote the subalgebra of $\mathfrak{g}(A)$ generated by the elements X_i (resp. Y_i), $i \in I$. Then one has the triangular decomposition $\mathfrak{g}(A) = \mathfrak{n}_- + \mathfrak{h} + \mathfrak{n}_+$.

A $\mathfrak{g}(A)$ -module V is called a *weight module* if $V = \oplus_{\mu \in \mathfrak{h}^*} V_\mu$, where $V_\mu = \{v \in V \mid hv = \mu(h)v, \text{ for all } h \in \mathfrak{h}\}$. If V_μ is non-zero, then μ is called a *weight*. A $\mathfrak{g}(A)$ -module V is called a *highest weight module* with *highest weight* $\lambda \in \mathfrak{h}^*$ if there exists a vector $v_\lambda \in V$ such that

$$\mathfrak{n}_+ v_\lambda = 0, \quad hv = \lambda(h)v_\lambda \text{ for } h \in \mathfrak{h}, \quad U(\mathfrak{g}(A))v_\lambda = V.$$

A highest weight module is a weight module. We let $L(\lambda)$ denote the irreducible highest weight module with highest weight λ .

A subalgebra $\mathfrak{s}$ of $\mathfrak{g}(A)$ is *locally finite* on a module V if $\dim U(\mathfrak{s})v < \infty$ for any $v \in V$. An element $x \in \mathfrak{g}$ is *locally nilpotent* on V if for any $v \in V$ there exists a positive integer N such that $x^N v = 0$.

For a Kac-Moody superalgebra $\mathfrak{g}(A)$ we do not necessarily have a non-degenerate symmetric invariant bilinear form, since we do not assume that the Cartan matrix A is symmetrizable. However, for a real non-isotropic root α and $\lambda \in \mathfrak{h}^*$ the quantity $\frac{2\lambda(h_\alpha)}{\alpha(h_\alpha)}$ is independent of the choice of $h_\alpha \in [\mathfrak{g}_\alpha, \mathfrak{g}_{-\alpha}]$, and we simply denote it by $\lambda(h_\alpha)$. If α is a real isotropic root, then $\lambda(h_\alpha)$ depends on the choice of h_α up to a non-zero scalar. To simplify notation we let $\lambda_i = \lambda(h_i)$, for $i \in I$.

Let $\mathfrak{g}$ be a finite-dimensional simple Lie superalgebra over $\mathbb{C}$ with a non-degenerate symmetric invariant bilinear form $(\cdot, \cdot)$. The associated (non-twisted) affine Lie superalgebra is

$$\widehat{\mathfrak{g}} = (\mathbb{C}[t, t^{-1}] \otimes_{\mathbb{C}} \mathfrak{g}) \oplus \mathbb{C}K \oplus \mathbb{C}d$$

with commutation relations

$$[a(m), b(n)] = [a, b](m+n) + m\delta_{m,-n}(a|b)K, \quad [d, a(m)] = -ma(m), \quad [K, \widehat{\mathfrak{g}}] = 0$$

where $a, b \in \mathfrak{g}$; $m, n \in \mathbb{Z}$ and $a(m) = t^m \otimes a$. By identifying $\mathfrak{g}$ with $1 \otimes \mathfrak{g}$, we have that the Cartan subalgebra of $\widehat{\mathfrak{g}}$ is

$$\mathfrak{h} = \overset{\circ}{\mathfrak{h}} \oplus \mathbb{C}K \oplus \mathbb{C}d$$

where $\overset{\circ}{\mathfrak{h}}$ is the Cartan subalgebra of $\mathfrak{g}$. Let δ be the linear function defined on $\mathfrak{h}$ by $\delta|_{\overset{\circ}{\mathfrak{h}} \oplus \mathbb{C}K} = 0$ and $\delta(d) = 1$. Let θ be the unique highest weight of the $\mathfrak{g}$. Then $\alpha_0 = \delta - \theta$ is the additional simple root which extends $\mathfrak{g}$ to $\widehat{\mathfrak{g}}$.

Now suppose σ is an automorphism of $\mathfrak{g}$ with order $m \neq 1$. Then $\mathfrak{g}$ decomposes into eigenspaces:

$$\mathfrak{g} = \bigoplus_{j \in \mathbb{Z}/m\mathbb{Z}} \mathfrak{g}(\sigma)_j, \quad \mathfrak{g}(\sigma)_j = \{x \in \mathfrak{g} \mid \sigma(x) = e^{\frac{j2\pi i}{m}} x\}.$$

The associated twisted affine Lie superalgebra is

$$\mathfrak{g}^{(m)} = \left(\bigoplus_{j \in \mathbb{Z}} t^j \otimes \mathfrak{g}(\sigma)_{j \bmod m} \right) \oplus \mathbb{C}K \oplus \mathbb{C}d$$

with the same commutation relations as given above. Let θ be the unique highest weight of the $\mathfrak{g}(\sigma)_0$ -module $\mathfrak{g}(\sigma)_1$. Then $\alpha_0 = \delta - \theta$ is the additional simple root which extends $\mathfrak{g}(\sigma)_0$ to $\mathfrak{g}^{(m)}$.

Let $\widehat{\mathfrak{L}}$ be an affine Lie algebra, and let V be a weight module over $\widehat{\mathfrak{L}}$. The module V is called an *integrable* module if for every real root α of $\widehat{\mathfrak{L}}$ the root space $\widehat{\mathfrak{L}}_\alpha$ is locally finite on V . Let $\widehat{\mathfrak{g}}$ be an affine Lie superalgebra, and let V

be a weight module over $\widehat{\mathfrak{g}}$. The module V is called an *integrable* module if it is integrable over the affine Lie algebra $\widehat{\mathfrak{g}}_0$. Let $V(\lambda)$ be a highest module over $\widehat{\mathfrak{g}}$. Then $k = \lambda(K)$ is the *level* of $L(\lambda)$.

It was shown in [9], that most non-twisted affine Lie superalgebras have only trivial irreducible integrable highest weight modules, and consequently the definition of integrable module was weakened.

Proposition 4.1. *[Kac, Wakimoto] The only non-twisted affine Lie superalgebras with non-trivial irreducible integrable highest weight modules are $osp(1, 2n)^{(1)}$, $C(n)^{(1)}$ and $A(0, m)^{(1)}$.*

In [2], S. Eswara Rao and V. Futorny show that over non-twisted affine Lie superalgebras all irreducible integrable weight modules with non-zero level are highest weight modules. The proof of the following statement is similar to the non-twisted case.

Proposition 4.2. *Let $\mathfrak{g}^{(m)}$ be a twisted affine Lie superalgebra which is not one of the algebras: $A(0, 2n - 1)^{(2)}$, $A(0, 2n)^{(4)}$ and $C(n)^{(2)}$. Then an irreducible integrable highest weight module over $\mathfrak{g}^{(m)}$ is trivial.*

Proof. First we consider the non-symmetrizable twisted affine Lie superalgebra, $q(n)^{(2)}$. If n is odd, then we have a Cartan matrix defined by $a_{i, i+1} = -1$, $a_{i+1, i} = 1$ for $i = 1, \dots, n \pmod{n}$, and all other entries zero. All simple roots in this case are odd isotropic. By an odd reflection at α_i , we obtain an even root $\alpha_i + \alpha_{i+1}$ with $h_{\alpha_i + \alpha_{i+1}} = h_{i+1} - h_i$. By Lemma 4.5, the conditions for integrability are $\lambda_{i+1} - \lambda_i \in \mathbb{Z}_{\geq 0}$, $(\text{mod } n)$. This implies $\lambda_i = 0$ for all $i = 1, \dots, n$.

If n is even, then we have a Cartan matrix defined by $a_{i, i+1} = -1$, $a_{i+1, i} = 1$ for $i = 1, \dots, n-1$, $a_{1, 1} = 2$, $a_{1, n} = a_{n, 1} = -1$, and all other entries zero. In this case, the simple root α_1 is even, and all other simple roots are odd isotropic. Using odd reflections, we obtain the set of principal roots $\{\alpha_1, \alpha_2 + \alpha_3, \alpha_3 + \alpha_4, \dots, \alpha_{n-1} + \alpha_n, \alpha_n + \alpha_1 + \alpha_2\}$. For $i = 2, \dots, n-1$ we have $h_{\alpha_i + \alpha_{i+1}} = h_{i+1} - h_i$, and $h_{\alpha_n + \alpha_1 + \alpha_2} = h_2 - h_n - h_1$. This yields integrability conditions $\lambda_1 \in \mathbb{Z}_{\geq 0}$, $\lambda_{i+1} - \lambda_i \in \mathbb{Z}_{\geq 0}$ for $i = 2, \dots, n-1$, and $\lambda_2 - \lambda_n - \lambda_1 \in \mathbb{Z}_{\geq 0}$. Together these imply $\lambda_i = 0$ for $i = 1, \dots, n$.

For a symmetrizable twisted affine Lie superalgebra $\mathfrak{g}^{(m)}$, we have the standard non-degenerate symmetric invariant bilinear form. The structure of $\mathfrak{g}_0^{(m)}$, the even part of $\mathfrak{g}^{(m)}$, is given by van de Leur in [12]. Let $\mathfrak{g}^{(m)}$ be a symmetrizable twisted affine Lie superalgebra, which is not one of the algebras: $A(0, 2n - 1)^{(2)}$, $A(0, 2n)^{(4)}$ and $C(n)^{(2)}$, and choose a base $\Pi = \{\alpha_0, \dots, \alpha_l\}$ with a unique simple isotropic root, α_d . Using the bilinear form to identify $\mathfrak{h}$ with $\mathfrak{h}^*$, we have $\alpha_i = \alpha_i^\vee$ for $i \leq d$ and $\alpha_i = -\alpha_i^\vee$ for $i > d$. Note that $d \neq 0$ and $d \neq l$ by choice of $\mathfrak{g}^{(m)}$.

Then $\mathfrak{g}_0^{(m)}$ has as subalgebras, both an affine subalgebra $\mathfrak{g}'$ on which this form is positive definite, and an affine subalgebra $\mathfrak{g}''$ on which this form is negative definite. Denote by $\overset{\circ}{\mathfrak{g}}'$ (resp. $\overset{\circ}{\mathfrak{g}}''$) the finite part of $\mathfrak{g}'$ (resp. $\mathfrak{g}''$). Let θ' (resp.

θ'') denote the highest weight of the module $\overset{\circ}{\mathfrak{g}}_1'$ (resp. $\overset{\circ}{\mathfrak{g}}_1''$) over $\overset{\circ}{\mathfrak{g}}_0'$ (resp. $\overset{\circ}{\mathfrak{g}}_0''$). Then the simple root $\alpha'_0 = \delta - \theta'$ (resp. $\alpha''_0 = \delta - \theta''$) extends $\overset{\circ}{\mathfrak{g}}_1'$ (resp. $\overset{\circ}{\mathfrak{g}}_1''$) to $\mathfrak{g}'$ (resp. $\mathfrak{g}''$). Also, $\theta' = \sum_{i=0}^{d-1} c_i \alpha_i$ and $\theta'' = \sum_{i=d+1}^l c_i \alpha_i$ with $c_i \in \mathbb{Z}_{>0}$.

Let $L(\lambda)$ be an irreducible integrable highest weight module over $\mathfrak{g}^{(m)}$. Let $k = \lambda(K)$, $k_i = \lambda(\alpha_i^\vee)$, $k' = \lambda(\alpha_0'^\vee)$ and $k'' = \lambda(\alpha_0''^\vee)$. Then by Lemma 4.5, $k_i \in \mathbb{Z}_{\geq 0}$ for $i \in I \setminus \{d\}$ and $k', k'' \in \mathbb{Z}_{\geq 0}$. Now using the bilinear form to identify $\mathfrak{h}$ with $\mathfrak{h}^*$, we have

$$\begin{aligned} \alpha_0'^\vee &= \alpha_0' &= \delta - \theta' &= K - \theta'^\vee \\ \alpha_0''^\vee &= -\alpha_0'' &= -(\delta - \theta'') &= -(K - \theta''^\vee). \end{aligned}$$

Hence,

$$\begin{aligned} k &= \lambda(K) &= \lambda(\alpha_0'^\vee) + \lambda(\theta'^\vee) &= k' + \lambda(\theta'^\vee) \\ -k &= -\lambda(K) &= \lambda(\alpha_0''^\vee) - \lambda(\theta''^\vee) &= k'' - \lambda(\theta''^\vee). \end{aligned}$$

Since $\theta'^\vee$ (resp. $\theta''^\vee$) is a positive (resp. negative) combination of $\alpha_i^\vee$ with $i \in I \setminus \{d\}$ and $\lambda(\alpha_i^\vee) \geq 0$ for $i \in I \setminus \{d\}$, we have that $\lambda(\theta'^\vee) \geq 0$ and $\lambda(\theta''^\vee) \leq 0$. Hence, $k = 0$ and the level of $L(\lambda)$ is zero. The above equations then imply that $k', k'', \lambda(\theta'^\vee), \lambda(\theta''^\vee) = 0$. Since $\lambda(\theta'^\vee) = \sum_{i=0}^{d-1} c_i k_i$ and $\lambda(\theta''^\vee) = -\sum_{i=d+1}^l c_i k_i$ with $c_i \in \mathbb{Z}_{>0}$, we have that $k_i = 0$ for $i \in I \setminus \{d\}$. Finally, $k' = 0$ then implies $k_d = 0$. Hence, the weight λ is zero, and so the module $L(\lambda)$ is trivial. $\square$

4.2 Integrable modules over regular Kac-Moody superalgebras

Let $\mathfrak{g}(A)$ be a regular Kac-Moody superalgebra, and let V be a weight module over $\mathfrak{g}(A)$. We call V *integrable* if for every real root α the element $X_\alpha \in \mathfrak{g}(A)_\alpha$ is locally nilpotent on V .

Lemma 4.3. *The adjoint module of a regular Kac-Moody superalgebra is an integrable module. In particular, ad_{X_α} is locally nilpotent for every real root α , where $X_\alpha \in \mathfrak{g}(A)_\alpha$.*

Proof. This follows from Lemma 1.3. $\square$

Our definition of integrable module coincides with the standard definition of integrable module in the case that $\mathfrak{g}(A)$ is an affine Lie superalgebra. This follows from the fact that every even root of an affine Lie superalgebra with non-zero length is real, as was shown in the dissertation of V. Serganova. The following lemma then follows immediately from Proposition 4.1 and Proposition 4.2.

Lemma 4.4. *Suppose $\mathfrak{g}(A)$ is a regular Kac-Moody superalgebra with a subfinite regular Kac-Moody diagram which is not of finite type and it is not one of the algebras: $A(0, m)^{(1)}$, $C(n)^{(1)}$, $S(1, 2, \alpha)$, and $Q^\pm(m, n, t)$. Then all irreducible integrable highest weight modules are trivial.*

The conditions for an irreducible highest weight module over $A(0, m)^{(1)}$ or $C(n)^{(1)}$ to be integrable were given in [9]. We are interested in the irreducible integrable highest weight modules for the regular Kac-Moody superalgebras: $S(1, 2, \alpha)$ and $Q^\pm(m, n, t)$. We will see that they have non-trivial irreducible integrable highest weight modules, and we will describe the weights.

Lemma 4.5. *Let $\mathfrak{g}(A)$ be a Kac-Moody superalgebra, and let $L(\lambda)$ be an irreducible highest weight module. If α_i is a simple non-isotropic root of $\mathfrak{g}(A)$, then $Y_i \in \mathfrak{g}(A)_{-\alpha_i}$ is locally nilpotent on $L(\lambda)$ if and only if $\lambda_i \in 2^{p(i)}\mathbb{Z}_{\geq 0}$. If α_i is a simple isotropic root, then $Y_i \in \mathfrak{g}(A)_{-\alpha_i}$ is locally nilpotent on $L(\lambda)$.*

Proof. If α_i is a simple even root, let $e = X_i$, $f = Y_i$ and $h = h_i$. Then $\{e, f, h\}$ is an s -triple. If α_i is a simple odd non-isotropic root, let $e = \frac{1}{4}[X_i, X_i]$, $f = \frac{1}{4}[Y_i, Y_i]$, and $h = \frac{1}{2}h_i$. Then $\{e, f, h\}$ is an s -triple. Since $\mathfrak{g}(A)$ is a Kac-Moody superalgebra, it follows that f is locally nilpotent on V if and only if f is nilpotent on the highest weight vector v_λ . Now f is nilpotent on v_λ if and only if $\lambda(h) \in \mathbb{Z}_{\geq 0}$. If α_i is odd, then $[Y_i, Y_i]v = 2(Y_i)^2v$ for $v \in V$. Thus, f is nilpotent on v_λ if and only if Y_i is nilpotent on v_λ . Hence, $Y_i \in \mathfrak{g}(A)_{-\alpha_i}$ is locally nilpotent on $L(\lambda)$ if and only if $\lambda_i \in 2^{p(i)}\mathbb{Z}_{\geq 0}$. If α_i is a simple isotropic root, then $(Y_i)^2v = 0$ for all $v \in L(\lambda)$. $\square$

Lemma 4.6. *Let $L(\lambda)$ be an irreducible integrable highest weight module of a regular Kac-Moody superalgebra $\mathfrak{g}(A)$, and let α_s be a simple root. Let $\mathfrak{n}'_+ = r_s(\mathfrak{n}_+)$ and $A' = r_s(A)$. Then after a simple even or odd reflection r_s the module $L(\lambda)$ is an irreducible integrable highest weight module of $\mathfrak{g}(A')$ with highest weight given below.*

1. If $\lambda(h_s) = 0$, then $v'_\lambda = v_\lambda$ is a highest weight vector with respect to $\mathfrak{n}'_+$ and the highest weight is $\lambda' = \lambda$.
2. If $\lambda(h_s) \neq 0$ and α_s is isotropic, then $v'_\lambda = Y_s v_\lambda$ is a highest weight vector with respect to $\mathfrak{n}'_+$ and the highest weight is $\lambda' = \lambda - \alpha_s$.
3. If $\lambda(h_s) = k \neq 0$ and α_s is even, then $v'_\lambda = (Y_s)^k v_\lambda$ is a highest weight vector with respect to $\mathfrak{n}'_+$ and the highest weight is $\lambda' = \lambda - k\alpha_s$.
4. If $\lambda(h_s) = 2n \neq 0$ and α_s is odd non-isotropic, then $v'_\lambda = (Y_s)^{2n} v_\lambda$ is a highest weight vector with respect to $\mathfrak{n}'_+$ and the highest weight is $\lambda' = \lambda - 2n\alpha_s$.

Proof. (1) and (2) follow immediately from the facts that $[Y_s, Y_s]v_\lambda = 2(Y_s)^2v_\lambda$, and that $Y_s v_\lambda = 0$ if and only if $\lambda(h_s) = 0$. For (3) and (4) suppose now that α_s is a non-isotropic root. Then by Lemma 4.5, we have $k = \lambda(h_s) \in 2^{p(s)}\mathbb{Z}_{\geq 0}$ since the module $L(\lambda)$ is assumed to be integrable. If α_s is a simple even root, then $\{X_s, Y_s, h_s\}$ is an s -triple. Set $v_j = \frac{1}{j!}(Y_s)^{(j)}v_\lambda$. Then $h_s v_j = (k - 2j)v_j$ and $X_s v_j = (k + 1 - j)v_{j-1}$. Thus $X_s v_{k+1} = 0$, and $X_i v_{k+1} = 0$ for all $i \in I \setminus \{s\}$. Since the module $L(\lambda)$ is irreducible this implies that $v_{k+1} = 0$. Thus $Y_s v_k = 0$ with $v_k \neq 0$. Hence, $L(\lambda)$ is a highest weight module with respect to $\mathfrak{n}'_+$, with highest weight vector $v_k = \frac{1}{k!}(Y_s)^k v_\lambda$ and the highest weight is $\lambda - k\alpha_s$.

Finally, suppose that α_s is an odd non-isotropic root. Set $v_j = (Y_s)^{(j)}v_\lambda$. Then $h_s v_j = (2n - 2j)v_j$, $X_s v_{2i} = (2i)v_{2i-1}$ and $X_s v_{2i-1} = (2n + 2 - 2i)v_{2i-2}$. Thus, $X_s(Y_s)^{2n+1}v_\lambda = 0$, and $X_i(Y_s)^{2n+1}v_\lambda = 0$ for $i \in I \setminus \{s\}$. Since the module $L(\lambda)$ is irreducible, we conclude that $(Y_s)^{2n+1}v_\lambda = 0$. Hence, $Y_s v_{2n} = 0$ with $v_{2n} \neq 0$. Therefore, $L(\lambda)$ is a highest weight module with respect to $\mathfrak{n}'_+$, with highest weight vector $(Y_s)^{2n}v_\lambda$ and highest weight $\lambda - (2n)\alpha_s$. The fact that the module $L(\lambda)$ is integrable as a $\mathfrak{g}(A')$ module follows from the fact that the real roots of $\mathfrak{g}(A)$ and $\mathfrak{g}(A')$ coincide. $\square$

Lemma 4.7. *Let $\mathfrak{g}(A)$ be a Kac-Moody superalgebra, and let $L(\lambda)$ be a highest weight module. Let α_s and α_i be simple non-isotropic roots. Let r_s be the even reflection with respect to α_s (or $2\alpha_s$ if α_s is odd). Suppose $Y_i \in \mathfrak{g}(A)_{-\alpha_i}$ is locally nilpotent on $L(\lambda)$, then $Y'_i \in \mathfrak{g}(A)_{-r_s(\alpha_i)}$ is locally nilpotent on $L(\lambda)$.*

Proof. The reflection r_s does not change the Cartan matrix, so $\mathfrak{g}(A')$ is again a Kac-Moody superalgebra. Thus, it suffices to show that Y'_i is nilpotent on the highest weight vector $v_{\lambda'}$ of $L(\lambda')$. It is sufficient to consider the case when both α_s and α_i are even roots. Then this is equivalent to the condition $\lambda'(h'_i) \in \mathbb{Z}_{\geq 0}$. Now $\lambda' = \lambda - \lambda(h_s)\alpha_s$. We have $\alpha'_s = -\alpha_s$ and $\alpha'_i = \alpha_i - a_{si}\alpha_s$. Also, $h'_s = -h_s$ and $h'_i = h_i - a_{is}h_s$. Then

$$\lambda'(h'_i) = (\lambda - \lambda(h_s)\alpha_s)(h_i - a_{is}h_s) = \lambda(h_i).$$

Since Y_i is locally nilpotent on $L(\lambda)$ we have $\lambda(h_i) \in \mathbb{Z}_{\geq 0}$. Hence Y'_i is locally nilpotent. $\square$

Corollary 4.8. *Let $\mathfrak{g}(A)$ be a regular Kac-Moody superalgebra. The irreducible highest weight module $L(\lambda)$ is an integrable module if and only if the element $Y_\alpha \in \mathfrak{g}(A)_{-\alpha}$ is locally nilpotent on $L(\lambda)$ for each principal root α .*

We call a weight λ *typical* if for any real odd root α we have $(\lambda + \rho)(h_\alpha) \neq 0$.

4.3 Integrable modules of the Lie superalgebra $S(1, 2, \alpha)$

Now we describe the weights for integrable highest weight modules of the Lie superalgebra $S(1, 2, \alpha)$. The Cartan matrix for the superalgebra $S(1, 2, \alpha)$ is

$$B = \begin{pmatrix} 2 & -1 & -1 \\ -1+b & 0 & 1 \\ -1-b & 1 & 0 \end{pmatrix}$$

with $b \neq \pm 1$.

Lemma 4.9. *Let $L(\lambda)$ be an irreducible highest weight module for $S(1, 2, \alpha)$. If $\lambda_2 \neq 0$ or $\lambda_3 \neq 0$, then $L(\lambda)$ is integrable if and only if*

$$\begin{aligned} \lambda_1 &\in \mathbb{Z}_{\geq 0} \\ \lambda_2 + \lambda_3 - 1 &\in \mathbb{Z}_{\geq 0}. \end{aligned}$$

If $\lambda_2 = \lambda_3 = 0$, then $L(\lambda)$ is integrable if and only if $\lambda_1 \in \mathbb{Z}_{\geq 0}$.

Proof. By Corollary 4.8, it suffices to find the conditions for the principal roots to be locally nilpotent on $L(\lambda)$. The principal roots of $S(1, 2, \alpha)$ are α_1 and $\alpha_2 + \alpha_3$. One can check that $h_{\alpha_2 + \alpha_3} = h_2 + h_3$. By Lemma 4.5, $Y_1 \in \mathfrak{g}(A)_{-\alpha_1}$ is locally nilpotent on $L(\lambda)$ if and only if $\lambda_1 \in \mathbb{Z}_{\geq 0}$. First suppose $\lambda_2 \neq 0$ and consider the odd reflection r_2 . We have that $r_2(\alpha_3) = \alpha_2 + \alpha_3$ and by Lemma 4.6 we have that $\lambda' = \lambda - \alpha_2$. Then by Lemma 4.5, $Y'_3 \in \mathfrak{g}(A)_{-(\alpha_2 + \alpha_3)}$ is locally nilpotent on $L(\lambda)$ if and only if $\lambda'(h'_3) \in \mathbb{Z}_{\geq 0}$ where

$$\lambda'(h'_3) = (\lambda - \alpha_2)(h_2 + h_3) = \lambda_2 + \lambda_3 - 1.$$

The argument for $\lambda_3 \neq 0$ is similar and yields the same condition. Finally, if $\lambda_2 = \lambda_3 = 0$ then the additional integrability condition is $\lambda(h_2 + h_3) \in \mathbb{Z}_{\geq 0}$, which is vacuously satisfied. $\square$

4.4 Integrable modules of the Lie superalgebra $Q^\pm(m, n, t)$

Now we describe the weights for integrable highest weight modules of the Lie superalgebra $Q^\pm(m, n, t)$. The Cartan matrix for $Q^\pm(m, n, t)$ is

$$\begin{pmatrix} 0 & 1 & a \\ b & 0 & 1 \\ 1 & c & 0 \end{pmatrix} \quad \begin{aligned} 1 + a + \frac{1}{b} &= m \\ 1 + b + \frac{1}{c} &= n \\ 1 + c + \frac{1}{a} &= t \end{aligned}$$

with $m, n, t \in \mathbb{Z}_{\leq -1}$, not all equal to -1, and the simple roots are odd isotropic. The principal even roots are $\{\alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_3\}$. One can check that for $i \neq j$,

$$h_{\alpha_i + \alpha_j} = \frac{h_j}{a_{ji}} + \frac{h_i}{a_{ij}}$$

and $(\alpha_i + \alpha_j)(h_{\alpha_i + \alpha_j}) = 2$.

Lemma 4.10. *A highest weight module $V(\lambda)$ for the algebra $Q^\pm(m, n, t)$ is typical if and only if $\lambda_1, \lambda_2, \lambda_3 \neq 0$.*

Proof. Since odd reflections of the diagram Γ do not yield new simple odd roots, the only conditions for the module to be typical are $\lambda(h_1), \lambda(h_2), \lambda(h_3) \neq 0$. $\square$

Lemma 4.11. *An irreducible highest weight module for $Q^\pm(m, n, t)$, with typical weight, is integrable if and only if*

$$\begin{aligned} \lambda_1 + \frac{1}{b}\lambda_2 - 1 \\ \lambda_2 + \frac{1}{c}\lambda_3 - 1 \\ \lambda_3 + \frac{1}{a}\lambda_1 - 1 \end{aligned} \in \mathbb{Z}_{\geq 0}.$$

Proof. By Corollary 4.8, it suffices to find the conditions for $Y_\alpha \in \mathfrak{g}(A)_{-\alpha}$ to be locally nilpotent on $L(\lambda)$ when α is a principal root. Let α_i be a simple isotropic root and let r_i be the odd reflection with respect to α_i . Since the weight λ is typical, $\lambda_i \neq 0$. Then by Lemma 4.6, $\lambda' = \lambda - \alpha_i$. Since the simple even roots of $\mathfrak{g}(A')$ are $\alpha_i + \alpha_j$ for $i \neq j$, we have by Lemma 4.5 that the conditions of integrability are $\lambda'(h'_j) \in \mathbb{Z}_{\geq 0}$, where

$$\lambda'(h'_j) = \lambda'(h_{\alpha_i + \alpha_j}) = (\lambda - \alpha_i) \left(\frac{h_j}{a_{ji}} + \frac{h_i}{a_{ij}} \right) = \frac{\lambda_j}{a_{ji}} + \frac{\lambda_i}{a_{ij}} - 1.$$

□

Proposition 4.12. *The non-trivial irreducible integrable highest weight modules of $Q^\pm(m, n, t)$ are $L(\lambda)$ such that*

$$\begin{pmatrix} \lambda_1 \\ \lambda_2 \\ \lambda_3 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 + abc \end{pmatrix} \begin{pmatrix} 1 & \frac{-1}{b} & \frac{1}{bc} \\ \frac{1}{ac} & 1 & \frac{-1}{c} \\ \frac{1}{a} & \frac{1}{ba} & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix},$$

with $x, y, z \in \mathbb{Z}_{>0}$. These weights are typical.

Proof. The dimension of the Cartan subalgebra is 3 by Corollary 3.2, and hence λ is determined by its values on h_1, h_2 and h_3 . First we consider the case when λ is typical. We can rewrite the conditions of Lemma 4.11 using matrices:

$$\begin{pmatrix} 1 & \frac{1}{b} & 0 \\ 0 & 1 & \frac{1}{c} \\ \frac{1}{a} & 0 & 1 \end{pmatrix} \begin{pmatrix} \lambda_1 \\ \lambda_2 \\ \lambda_3 \end{pmatrix} = \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

with $x, y, z \in \mathbb{Z}_{>0}$. The determinant of the left most matrix is $\frac{1+abc}{abc}$, which is nonzero by Corollary 3.2. Hence, the matrix is invertible, and is calculated above. Finally, suppose that λ is not typical. Then without loss of generality suppose $\lambda_1 = 0$. Consider the odd reflection r_1 with respect to α_1 . By Lemma 4.6, $\lambda' = \lambda$. Then by Lemma 4.5, we have the integrability conditions $x, z \in \mathbb{Z}_{\geq 0}$ with

$$\begin{aligned} x &= \lambda'(h_{\alpha_1 + \alpha_2}) = \lambda_1 + \frac{1}{b}\lambda_2 = \frac{1}{b}\lambda_2 \\ z &= \lambda'(h_{\alpha_1 + \alpha_3}) = \lambda_3 + \frac{1}{a}\lambda_1 = \lambda_3. \end{aligned}$$

If λ_2 or λ_3 is nonzero, then by a reflection at the corresponding simple root we obtain the integrability condition $y \in \mathbb{Z}_{\geq 0}$ with

$$y = \lambda_2 + \frac{1}{c}\lambda_3 - 1.$$

By Lemma 3.1, $b, c < 0$. But this together with $\frac{1}{b}\lambda_2, \lambda_3 \geq 0$ implies $\lambda_2 + \frac{1}{c}\lambda_3 - 1 < 0$, which is a contradiction. Hence, if λ is not typical then $\lambda = 0$. □

5 Extending regular Kac-Moody diagrams that are not of finite type

In this section, we prove that a subfinite regular Kac-Moody diagram with an isotropic vertex, which is not of finite type, is not extendable. Hence, a regular Kac-Moody diagram with an isotropic vertex is subfinite.

Lemma 5.1. *Suppose Γ and $\Gamma' = \Gamma \cup \{v_{n+1}\}$ are connected regular Kac-Moody diagrams. Let $\mathfrak{g}(A)$ (resp. $\mathfrak{g}(A')$) be the Kac-Moody superalgebra with diagram Γ (resp. Γ'), and let $Y_{n+1} \in \mathfrak{g}(A')_{-\alpha_{n+1}}$ be the generator corresponding to the vertex v_{n+1} . Then the submodule M of $\mathfrak{g}(A')$ generated by $\mathfrak{g}(A)$ acting on Y_{n+1} is an integrable highest weight module over the subalgebra $\mathfrak{g}(A)$.*

Proof. The fact that M is a highest weight module follows immediately from $[X_i, Y_{n+1}] = 0$ for all $i = 1, \dots, n$. The module M has highest weight $-\alpha_{n+1}$. A real root α of the subalgebra $\mathfrak{g}(A)$ is also a real root of $\mathfrak{g}(A')$. By Lemma 4.3, the adjoint module of $\mathfrak{g}(A')$ is integrable. Thus for each real root α of $\mathfrak{g}(A)$ we have that $Y_\alpha \in \mathfrak{g}(A)_{-\alpha}$ acts locally nilpotently on the submodule M of $\mathfrak{g}(A')$. Hence the submodule M is an integrable highest weight module over the subalgebra $\mathfrak{g}(A)$. $\square$

Corollary 5.2. *If Γ is a diagram for a regular Kac-Moody superalgebra $\mathfrak{g}(A)$ that does not have non-trivial irreducible integrable highest weight modules, then Γ is not extendable.*

Proof. If $\mathfrak{g}(A)$ has only trivial irreducible integrable highest weight modules, then the highest weight of the module M is 0. Hence $-\alpha_{n+1} = 0$, which implies $a_{i,n+1} = 0$ for $i = 1, \dots, n$. Since we assumed that the matrix A is a generalized Cartan matrix, this implies $a_{n+1,j} = 0$ for $j = 1, \dots, n$. This is not possible with Γ' being a connected diagram. $\square$

Corollary 5.3. *Suppose that Γ is a subfinite regular Kac-Moody diagram for $\mathfrak{g}(A)$ which is not of finite type and not one of the algebras: $A(0, m)^{(1)}$, $C(n)^{(1)}$, $S(1, 2, \alpha)$, and $Q^\pm(m, n, t)$. Then the diagram Γ is not extendable.*

Proof. This follows immediately from Lemma 4.4. $\square$

The following observation will be used in the proof of the next theorem. It follows from the classification of 3-vertex regular Kac-Moody diagrams.

Remark 5.4. If Γ is a regular Kac-Moody 3-vertex diagram which is not a diagram for $Q^\pm(m, n, t)$, $S(1, 2, \alpha)$ or $D(2, 1, \alpha)$, then the ratio of an isotropic vertex in Γ is rational and negative. The ratio of an isotropic vertex for $Q^\pm(m, n, t)$ is real, irrational and negative. For $S(1, 2, \alpha)$ and $D(2, 1, \alpha)$, if the ratios are rational then at most one of them is positive.


Theorem 5.5. *A connected regular Kac-Moody diagram containing an isotropic vertex is subfinite. In particular, if Γ is a subfinite regular Kac-Moody diagram which is not of finite type, then Γ is not extendable.*

Proof. We prove this by induction on the number of vertices. This is clearly true for a one vertex diagram. Suppose that the claim is true for all diagrams with less than n vertices, and let Γ be a connected n -vertex diagram which is regular Kac-Moody and contains an isotropic vertex. If Γ' is a connected proper subdiagram of Γ containing an isotropic vertex, then Γ' is regular Kac-Moody and by the induction hypothesis Γ' is subfinite.

Let $\mathcal{S}$ denote the collection of connected subfinite regular Kac-Moody which are extendable. Then $\Gamma' \in \mathcal{S}$ is either of finite type or a diagram for $A(0, m)^{(1)}$, $C(n)^{(1)}$, $S(1, 2, \alpha)$, $Q^\pm(m, n, t)$. This follows from the classification of subfinite regular Kac-Moody diagrams and Corollary 5.3.

In the following lemmas, we prove that if Γ' is a diagram for: $A(0, m)^{(1)}$, $C(n)^{(1)}$, $S(1, 2, \alpha)$, or $Q^\pm(m, n, t)$, then Γ' is not a proper subdiagram of a connected regular Kac-Moody diagram Γ which satisfies the condition: all proper connected regular Kac-Moody diagrams containing an isotropic vertex are in $\mathcal{S}$.

Lemma 5.6. $C(n)^{(1)} \notin \mathcal{S}$




Proof. Let v_{n+2} denote the additional vertex. Now $\Gamma_1 \in \mathcal{S}$ implies $a_{n+2, n+1}, a_{n+2, n} = 0$, and $\Gamma_{n+1} \in \mathcal{S}$ implies $a_{n+2, j} = 0$ for $j = 1, \dots, n-1$. Hence, $C(n)^{(1)}$ is not extendable. $\square$

Lemma 5.7. $Q^\pm(m, n, t) \notin \mathcal{S}$

Proof. We consider each case for attaching a vertex to a $Q^\pm(m, n, t)$ diagram. Let Γ denote the corresponding extended diagram. Recall that $m, n, t \in \mathbb{Z}_{<0}$ and are not all equal to -1 , and that $a, b, c \in (\mathbb{R} \setminus \mathbb{Q})_{<0}$ satisfy


$$\begin{aligned} 1 + a + \frac{1}{b} &= m \\ 1 + b + \frac{1}{c} &= n \\ 1 + c + \frac{1}{a} &= t. \end{aligned} \tag{11}$$

Case 1:




By Remark 5.4, we have that $d, \frac{b}{d} \in \mathbb{Q}$, implying $b \in \mathbb{Q}$, which is a contradiction.

Case 2:



Now $\Gamma_1 \in \mathcal{S}$ implies $g < 0$, and $\Gamma_3 \in \mathcal{S}$ implies $\frac{e}{d} < 0$. Since $a, b, c < 0$, this implies $d, \frac{g}{a} > 0$. By Remark 5.4, this implies $\Gamma_2 \notin \mathcal{S}$, which is a contradiction.

Case 3:



The ratios of v_1, v_2, v_3 in Γ_4 are $a, b, c \in (\mathbb{R} \setminus \mathbb{Q})_{<0}$. If Γ_1 is a subdiagram such that the ratios of v_2 and v_3 in Γ_1 are real negative numbers. Then the ratio of v_2 in Γ_3 and of v_3 in Γ_2 are real positive numbers. But then by Remark 5.4 all of the ratios of v_1 are real negative numbers, which is a contradiction. Hence, $\Gamma_1, \Gamma_2, \Gamma_3$ are diagrams for $D(2, 1, \alpha)$ or $S(1, 2, \alpha)$.

Subcase 1: v_4 is $\otimes$ and $\Gamma_1, \Gamma_2, \Gamma_3$ are diagrams for $D(2, 1, \alpha)$.

Then by (2) the variables satisfy:


$$\begin{array}{lll} h + \frac{g}{a} = -1 & \frac{c}{h} + \frac{d}{k} = -1 & g + \frac{e}{b} = -1 \\ \frac{a}{g} + \frac{k}{f} = -1 & \frac{k}{d} + \frac{1}{e} = -1 & \frac{b}{e} + \frac{f}{d} = -1 \\ \frac{f}{k} + \frac{1}{h} = -1 & e + \frac{h}{c} = -1 & \frac{d}{f} + \frac{1}{g} = -1 \end{array}.$$

Now solving for h we find $h = \frac{bc-c-abc}{1+abc} \in \mathbb{R}$. Similarly, we find that $d, e, f, g, k \in \mathbb{R}$ and all ratios are real. Since one ratio at each vertex v_1, v_2, v_3 must be positive, and the diagram for $D(2, 1, \alpha)$ has at most one positive ratio vertex, it follows that all of the ratios at v_4 are negative, which is a contradiction.

Subcase 2: v_4 is $\circ$ and $\Gamma_1, \Gamma_2, \Gamma_3$ are diagrams for $S(1, 2, \alpha)$.

Then $d, f, k = -1$, $h + \frac{g}{a} = -2$, $g + \frac{e}{b} = -2$, and $e + \frac{h}{c} = -2$. Solving for e we find $e = \frac{2(ab-b-abc)}{1+abc} \in \mathbb{R}$. Without loss of generality, suppose $a + \frac{1}{b} + 1 \in \mathbb{Z}_{\leq -2}$.

Now by reflecting at v_2 we have that Γ' is



which implies that Γ'_2 is a diagram $Q^\pm(m, n, t)$. Then the ratio of v_4 in Γ'_2 is $\frac{-b-2e}{-1-2e} \in (\mathbb{R} \setminus \mathbb{Q})_{<0}$. But by substituting $e = \frac{2(ab-b-abc)}{1+abc}$ we have


$$\frac{-b-2e}{-1-2e} = \frac{3b-ab^2c-4ab+4abc}{-1+3abc-4ab+4b} > 0,$$

which is a contradiction. Therefore, $Q^\pm(m, n, t)$ is not extendable. $\square$


Lemma 5.8. $S(1, 2, \alpha) \notin \mathcal{S}$

Proof. Note that an odd reflection of a $S(1, 2, \alpha)$ diagram is again a $S(1, 2, \alpha)$ diagram, but with a different α .

Case 1:




◊If v_4 is $\otimes$, then $\Gamma_2 \in \mathcal{S}$ implies $c = 1$ and $b = a$. By reflecting at v_4 we have



But $\Gamma'_4 \notin \mathcal{S}$.


◊If v_4 is $\odot$ and $c = -2$, then $\Gamma_2 \in \mathcal{S}$ implies $b = a$ and $\Gamma_3 \in \mathcal{S}$ implies

$\frac{a-1}{a} \in \{-1, -2, \frac{-3}{2}\}$. Hence, $\frac{1}{3} \leq a \leq \frac{1}{2}$. By reflecting at v_3 and then at v_2 , we obtain $\Gamma'' := r_2(r_3(\Gamma))$




Then $\Gamma''_2 \in \mathcal{S}$ implies $\frac{a+3}{a+2} \in \{-1, -2, \frac{-3}{2}\}$, which is a contradiction.

◊ If v_4 is $\bigcirc$ and $c = -1$, then $\Gamma_2 \in \mathcal{S}$ implies $\frac{b}{a} \in \{1, 2, 3\}$. Reflecting at v_1 yields Γ'




First suppose that $b = 1-a$. Then for $\Gamma_1 \in \mathcal{S}$ implies $\frac{a-b}{a} \in \{0, -1\}$. If $\frac{a-b}{a} = 0$, then we are reduced to a previous case. If $\frac{a-b}{a} = -1$, then $a = \frac{1}{3}$. Reflection at v_4 of Γ' yields Γ''




But, $\Gamma''_4 \notin \mathcal{S}$.

Now we assume that $b \neq 1 - a$. Then reflecting at v_2 of Γ' yields Γ''




Now $\Gamma''_2 \in \mathcal{S}$ implies $\frac{3-a-2b}{1-a-b} \in \{0, -1, -2\}$. Recall that $a \notin \mathbb{Z}$ and $\frac{b}{a} \in \{1, 2, 3\}$. If $\frac{3-a-2b}{1-a-b} = 0$, then $a + 2b - 1 - \frac{2b}{a} = 0$. This implies $a = 1$ or $\frac{b}{a} = \frac{-1}{2}$, which is a contradiction. If $\frac{3-a-2b}{1-a-b} = -2$, then Γ''_1 is a $G(3)$ diagram and $a = b = \frac{5}{7}$. But then $\Gamma''_2 \notin \mathcal{S}$. If $\frac{3-a-2b}{1-a-b} = -1$, then by solving for b we have $b = \frac{4-2a}{3}$. Now $\Gamma''_1 \in \mathcal{S}$ implies $\frac{1-a-b}{-a+2} \in \{-1, \frac{-1}{2}\}$. Solving these equations yields $a, b = \frac{4}{5}$ and Γ''_1 is a diagram for $C(3)$. By substituting and then reflecting Γ'' at v_3 we obtain Γ'''




But then $\Gamma'''_3 \notin \mathcal{S}$.

Case 2:




By reflecting at v_1 , we are reduced to Case 1.

Case 3:




Now $\Gamma_2 \in \mathcal{S}$ implies v_4 is either $\otimes$ or $\circ$.

◊If v_4 is $\otimes$, then $\Gamma_2 \in \mathcal{S}$ implies $b = c, d = e$, and $e = a - c$. Also, $\Gamma_1, \Gamma_3 \in \mathcal{S}$ implies $\frac{a-1}{c}, \frac{a+1}{a-c} \in \{-1, -2, -3\}$. If $\frac{a-1}{c} = \frac{a+1}{a-c}$, then $c = \frac{a-1}{2}$ or $\frac{a-1}{c} = 2$, which is a contradiction. If either fraction equals -1 , then a reflection at the corresponding vertex returns us to Case 1. So without loss of generality by symmetry we have that $\frac{a-1}{c} = -2$ and $\frac{a+1}{a-c} = -3$. Then $a = \frac{1}{11}$ and $c = \frac{5}{11}$. By substituting and then reflecting at v_3 , we obtain Γ'




But, $\Gamma'_3 \notin \mathcal{S}$.

◊If v_4 is $\circ$, then $\Gamma_2 \in \mathcal{S}$ implies that either $b, d = -1$ or $b = -1, d = -2$, without loss of generality by symmetry. First suppose that $b = -1, d = -2$. Then Γ_2 is a $G(3)$ diagram and $c = \frac{a}{3}, e = \frac{2a}{3}$. Then $\Gamma_1 \in \mathcal{S}$ implies $e = -a - 1$. Thus $a = -\frac{3}{5}$. By substituting and then reflecting at v_1 , we obtain Γ'




But, $\Gamma'_1 \notin \mathcal{S}$. Now assume that $b = d = -1$. Then $\Gamma_2 \in \mathcal{S}$ implies that either

$-\frac{c}{a} = -\frac{e}{a} = -\frac{1}{2}$ or $-\frac{c}{a} - \frac{e}{a} = -2$. Now by reflecting Γ at v_1 we obtain Γ'



If $-\frac{c}{a} = -\frac{e}{a} = -\frac{1}{2}$, then Γ' reduces to a previous case, so we assume not. Thus, $c+e=2a$ and $\frac{a-e-c}{a} = -1$. If $1-a-c \neq 0$, then $\Gamma'_1 \in \mathcal{S}$ implies that either $\frac{a-2c}{1-a-c} = \frac{2-a}{1-a-c} = -\frac{1}{2}$ or $\frac{a-2c}{1-a-c} + \frac{2-a}{1-a-c} = -2$. Now by reflecting Γ at v_3 we obtain Γ''




If $-1-a-e \neq 0$, then $\Gamma''_3 \in \mathcal{S}$ implies that either $\frac{-a-2}{-1-a-e} = \frac{a-2e}{-1-a-e} - \frac{1}{2}$, or $\frac{-a-2}{-1-a-e} + \frac{a-2e}{-1-a-e} = -2$.

If $c = 1-a$, then $e = 3a-1$ and $-1-a-e = -4a \neq 0$. Then by substitution we have that $\Gamma''_3 \in \mathcal{S}$ implies that either $\frac{-a-2}{-4a} = \frac{2-5a}{-4a} = -\frac{1}{2}$, or $\frac{-a-2}{-4a} + \frac{2-5a}{-4a} = -2$, but since $a \notin \mathbb{Z}$ these equations have no solution. Hence, $1-a-c \neq 0$.

If $e = -1-a$, then $c = 3a+1$. Then by substitution we have that $\Gamma'_1 \in \mathcal{S}$ implies that either $\frac{-2-5a}{-4a} = \frac{2-a}{-4a} = -\frac{1}{2}$ or $\frac{-2-5a}{-4a} + \frac{2-a}{-4a} = -2$, but since $a \notin \mathbb{Z}$ these equations have no solution. Hence, $-1-a-e \neq 0$.


If $\frac{a-2c}{1-a-c} = \frac{2-a}{1-a-c} = -\frac{1}{2}$, then reflecting Γ' at v_4 reduces us to Case 1. Thus $\frac{a-2c}{1-a-c} + \frac{2-a}{1-a-c} = -2$, which implies $c = 1 - \frac{a}{2}$. If $\frac{-a-2}{-1-a-e} = \frac{a-2e}{-1-a-e} = -\frac{1}{2}$, then reflecting Γ'' at v_4 reduces us to Case 1. Thus $\frac{-a-2}{-1-a-e} + \frac{a-2e}{-1-a-e} = -2$, which implies $e = -1 - \frac{a}{2}$. But then $c+e = -a$, which is a contradicts $c+e = 2a$.

Case 4:



Now $\Gamma_1 \in \mathcal{S}$ implies v_4 is $\otimes$ and $\Gamma_2 \in \mathcal{S}$ implies $e = a$. By reflecting at v_3 , we are reduced to Case 3.

Case 5:




If v_4 is $\bigcirc$, then $\Gamma_1 \notin \mathcal{S}$. If v_4 is $\otimes$ then Γ_2 is $D(2, 1, \alpha)$. But then a reflection at v_1 reduces to a previous case. Therefore, $S(1, 2, \alpha)$ is not extendable. $\square$

Lemma 5.9. $A(0, m)^{(1)} \notin \mathcal{S}$


Proof. First we show that $A(0, 1)^{(1)}$ is not extendable.

Case 1:




Now $\Gamma_2 \in \mathcal{S}$ implies $a < 0$. Then $\Gamma_3 \in \mathcal{S}$ implies Γ_3 is a diagram for $D(2, 1, \alpha)$ diagram, where v_4 is $\bigcirc$ and $b = -1$. Then $\Gamma_2 \in \mathcal{S}$ implies $c \in \{-1, -2, -3\}$.

By reflecting at v_1 we obtain Γ'




Then $\Gamma'_1 \notin \mathcal{S}$ since $1 - c > 0$.

Case 2:




By reflecting at v_1 we return to Case 1.

Case 3:




Now $\Gamma_2 \in \mathcal{S}$ implies $e < 0$. Then $\Gamma_3 \notin \mathcal{S}$.

Case 4:




◊If v_4 is $\otimes$ then $\Gamma_1, \Gamma_3 \in \mathcal{S}$ implies that $e, c \in \mathbb{Q}_{>0}$. But then $\Gamma_2 \notin \mathcal{S}$ since it has two vertices with positive ratios.

◊If v_4 is $\circ$, then Γ_1 and Γ_3 are $D(2, 1, \alpha)$ diagrams. So $b, d = -1$. Then $\Gamma_2 \in \mathcal{S}$ implies $e = c \in \{-1, -\frac{1}{2}\}$. By reflecting at v_1 we obtain Γ'




If $c = -\frac{1}{2}$, then this is a previous case. If $c = -1$, then $\Gamma'_1 \notin \mathcal{S}$.

Case 5:



Now $\Gamma_3 \in \mathcal{S}$ implies $c > 0$. Then $\Gamma_2 \in \mathcal{S}$ implies that Γ_2 is a diagram for $D(2, 1, \alpha)$. So v_4 is $\otimes$, $b = c$, and $d = e = -1 - c$. But then $\Gamma_3 \notin \mathcal{S}$ since the ratio at v_3 is positive. Therefore, $A(0, 1)^{(1)}$ is not extendable.

Now we show that $A(0, 2)^{(1)}$ is not extendable.




First suppose that $a_{15} \neq 0$.

◊If v_5 is $\otimes$ then $\Gamma_{3,4}$ is $D(2, 1, \alpha)$, and so $a_{2,5} \neq 0$. Then $\Gamma_4 \in \mathcal{S}$ implies $a_{3,5} = 0$, and $\Gamma_3 \in \mathcal{S}$ implies $a_{4,5} = 0$. Then $\Gamma_{1,4} \in \mathcal{S}$ implies $a_{25} > 0$, and $\Gamma_{2,3} \in \mathcal{S}$ implies $a_{15} > 0$. Then $\Gamma_{3,4}$ has two isotropic vertices with positive ratio, which is a contradiction.

◊If v_5 is $\circ$ then $\Gamma_{2,3}$ is $D(2, 1, \alpha)$, and so $a_{51} = -1$. Then by reflecting at v_1 we return a previous case.

Therefore, $a_{15} = 0$ and by symmetry $a_{25} = 0$. Now without loss of generality $a_{45} \neq 0$. Then by reflecting at v_1 we have v'_4 is $\otimes$ and $a'_{45} \neq 0$, but this is the previous case. Hence, $A(0, 2)^{(1)}$ is not extendable.

Finally we show that $A(0, m)^{(1)}$ is not extendable, for $m \geq 3$. Let Γ be a diagram for $A(0, m)^{(1)}$. Then Γ has $m + 2 \geq 5$ vertices. Let v_1 denote the vertex being added to the diagram. First suppose v_1 is connected to an isotropic vertex, which we denote v_2 . Let v_3 and v_4 denote the vertices adjacent to v_2 in Γ' . Since v_2 is isotropic with degree 3, it must be contained in a $D(2, 1, \alpha)$ subdiagram. Since the subdiagram $\{v_3, v_2, v_4\}$ is not $D(2, 1, \alpha)$, we have without loss of generality that the subdiagram $\{v_1, v_2, v_3\}$ is $D(2, 1, \alpha)$. Then v_1 is either $\otimes$ or $\circ$. If v_1 is $\otimes$ then we have the following subdiagram:



where the double lines are necessarily connected, and the dotted lines are possibly connected. By the 5-vertex classification, this diagram is not subfinite regular Kac-Moody. If v_1 is $\circ$, then by reflecting at v_2 we return to the this case.

Next suppose v_1 is not connected to an isotropic vertex. Then let Γ'' be a minimal subdiagram containing v_1 and an isotropic vertex, which we denote v_n . By minimality of Γ'' it is a chain such that v_i is not isotropic for $1 < i < n$. Thus by Lemma 2.4, there is a sequence of odd reflections R such that $R(v_2)$ is isotropic and connected to v_1 . This reduces us to the previous case. $\square$

This concludes the proof of Theorem 5.5. $\square$

Remark 5.10. Theorem 2.26 follows from the classification of connected subfinite regular Kac-Moody diagrams in Section 2 and from Theorem 5.5.

References

- [1] V.K. Dobrev and V.B. Petkova, *On the group-theoretical approach to extended conformal supersymmetry: function space realizations and invariant differential operators*, Fortschr. d. Phys. **35** (1987) 537-572.
- [2] S. Eswara Rao and V. Futorny, *Irreducible modules for affine Lie superalgebras*, e-print: math/0311207 (2003).
- [3] C. Hoyt, *Kac-Moody superalgebras of finite growth*, Ph.D. thesis, University of California, Berkeley (2007).
- [4] C. Hoyt and V. Serganova, *Classification of finite-growth general Kac-Moody superalgebras*, Communications in Algebra **35 no. 3** (2007), 851-874.
- [5] V.G. Kac, *Lie superalgebras*, Advances in Math **26** (1977), 8-96.
- [6] V.G. Kac, *Infinite-dimensional algebras, Dedekind's η -function, classical Möbius function and the very strange formula*, Advances in Math **30** (1978), 85-136.
- [7] V.G. Kac, *Infinite dimensional Lie algebras, 3rd ed.*, Cambridge University Press, 1990.
- [8] V.G. Kac and J.W. van de Leur, *On classification of superconformal algebras*, Strings **88**, World Sci. (1989), 77-106.
- [9] V.G. Kac and M. Wakimoto, *Integrable highest weight modules over affine superalgebras and Appell's function*, Comm. Math. Phys. **215** (2001), no. 3, 631-682.
- [10] D. Leites, M. Savel'ev and V. Serganova, *Embeddings of Lie superalgebra $osp(1|2)$ and nonlinear supersymmetric equations*, Group Theoretical Methods in Physics **1** (1986), 377-394.
- [11] V. Serganova, *Kac-Moody superalgebras and integrability*, Perspectives and trends in infinite-dimensional Lie theory.
- [12] J.W. van de Leur, *Contragredient Lie superalgebras of finite growth*, Thesis Utrecht University (1986).
- [13] J.W. van de Leur, *A classification of contragredient Lie superalgebras of finite growth*, Communications in Algebra **17** (1989), 1815-1841.